AUTHOR INDEX

(D) Denotes Demonstration, (T) Title only, (C) Communication, (PC) Poster Communication

ABBOTT N. Joan & BROWN E.R. Effects of osmotic manipulation on extracellular routes for ion regulation in the squid Schwann cell. (C) 16*P*

ACATAUASSÚ-NUNES R., ALBUQUERQUE A.M.T., PRIST R. & ROCHA E SILVA M. Haemodynamics of cardiac tamponade after hypertonic NaCl (7·5%) Dextran-70 (6%) – HSD – injections in anaesthetized dogs. (C) 81*P*

ADAMS D.J., VAN BREEMEN C., CANNELL M.B. & SAGE S.O. Na:Ca exchange in Fura-2-loaded bovine pulmonary artery endothelial cell monolayers. (PC) 183*P*

ADAMS M.E., BINDOKAS V.P., DOLPHIN A.C. & SCOTT R.H. The inhibition of Ca^{2+} channel currents in cultured neonatal rat dorsal root ganglion (DRG) neurones by ω-Agatoxin-1A (a funnel web spider toxin). (C) 34*P*

AGUS Z.S., DUKES I.D. & MORAD M. Divalent cations modulate transient outward current in isolated rat ventricular myocytes. (C) 28*P*

AIKEN S.P., HORN N.M. & SAUNDERS N.R. Effect of histidine on zinc distribution in anaesthetized rats. (C) 144*P*

AL-BARAZANJI K.A. & BALMENT R.J. Central and peripheral effects of the peptide ANF on renal function and blood pressure in hypertensive rats. 261

ALBUQUERQUE A.M.T. *see* ACATAUASSÚ-NUNES R., ALBUQUERQUE A.M.T., PRIST R. & ROCHA E SILVA M. Haemodynamics of cardiac tamponade after hypertonic NaCl (7·5%) Dextran-70 (6%) – HSD – injections in anaesthetized dogs. (C) 81*P*

ALLEN T.StC. & GORDON A.M. Cycling cross-bridges produce structural changes in Troponin-C in demembranated fibres of rabbit psoas muscle. (C) 63*P*

ALMERS W., IWATA A., SPRUCE A.E. & WHITE J.M. Patch clamp studies of fusion of human erythrocytes to murine fibroblasts mediated by influenza virus fusion protein. (C) 95*P*

ANDERSON L.C. & SULEIMAN A.H. Effects of streptozotocin-induced diabetes on sympathetically evoked submandibular salivary secretion in anaesthetized rats. (C) 98*P*

ANDERSON L.S., BUTLER P.J., ROBERTS C., SMALE K., SNOW D.H. & WOAKES A.J. Partial pressures of arterial and end-tidal gases during graded exercise in thoroughbred race-horses. (C) 131*P*

ANDERSSON P.-O., SJÖGREN C., UVNÄS B. & UVNÄS-MOBERG K. The role of vasoactive intestinal polypeptide (VIP) in mediating responses of the urinary bladder to stimulation of the pelvic and hypogastric nerves in anaesthetized dogs. (C) 74*P*

AOTA S. & RANDALL D.J. Catecholamines and the central control of breathing in fish. (C) 130*P*

ASHLEY C.C., BAGNI A., CECCHI G., GRIFFITHS P.J. & MAEDA Y. Time-resolved X-ray diffraction studies on intact single muscle fibres of *Rana temporaria*. (C) 58*P*

BACKHOUSE S.S., HARPER A.A., VAN VLIET B.N. & WEST N.H. Localization and conduction velocities of arterial baroreceptors in the anaesthetized snake. (C) 135*P*

BAGNI A. *see* ASHLEY C.C., BAGNI A., CECCHI G., GRIFFITHS P.J. & MAEDA Y. Time-resolved X-ray diffraction studies on intact single muscle fibres of *Rana temporaria*. (C) 58*P*

BAHINSKI Anthony, GADSBY David C., GREENGARD Paul & NAIRN Angus C. Chloride conductance regulated by protein kinase A in isolated guinea-pig ventricular myocytes. (C) 32*P*

BAKER G.F. & WIDDAS W.F. Red cell volume compensation in isosmotic glucose. (C) 87*P*

BALKE C.W. & WIER W.G. Two types of calcium current in isolated guinea-pig ventricular myocytes: accurate separation by voltage-dependence of steady-state inactivation. (C) 31*P*

BALL F.G., SANSOM M.S.P. & USHERWOOD P.N.R. A comparison of different gating models for voltage- and receptor-gated ion channels. (PC) 189*P*

BALLESTA J.J., BORGES R., GARCÍA A.G. & HIDALGO M.J. Secretory and radioligand binding studies on muscarinic receptors in bovine and feline chromaffin cells. 411

BALMENT R.J. *see* AL-BARAZANJI K.A. & BALMENT R.J. Central and peripheral effects of the peptide ANF on renal function and blood pressure in hypertensive rats. 261

BARKER P.M., MARKIEWICZ M., PARKER K.A., STRANG L.B. & WALTERS D.V. Induction of the adrenaline-dependent reabsorption of lung liquid in the fetal sheep by synergistic action of triiodothyronine and hydrocortisone. (C) 138*P*

BARKLEY G.L., GARDNER-MEDWIN A.R., MORAN J., NAGEL-LEIBY S., SIMKINS R.T., TEPLEY N. & WELCH K.M.A. Magnetic fields during spreading depression in the cortex of anaesthetized rabbits. (C) 101*P*

BARO I., EISNER D.A., FRY C.H., LISTON T.G., MONTGOMERY B.S.I. & RAIMBACH S.J. Measurement of intracellular pH in isolated ferret detrusor smooth muscle cells. (C) 124*P*

BARROS L.F., BUSTAMANTE J.C., JARVIS S.M. & YUDILEVICH D.L. Binding of nitrobenzylthioinosine, an inhibitor of facilitated-diffusion nucleoside transport, to brush-border and basal membrane vesicles isolated from the human placenta. (PC) 179*P*

BASKIN R.J., SHEN S. & YEH Y. Optical measurements on isolated single fibres from insect flight muscle. (C) 55*P*

BAYLOR S.M., HOLLINGWORTH S. & KONISHI M. Calcium transients in intact single frog skeletal muscle fibres measured with the fluorescence indicator dye Mag-fura-2. (C) 69*P*

BEECH D.J. & BOLTON T.B. Two components of potassium current activated by depolarization of single smooth muscle cells from the rabbit portal vein. 293

BEN ARI Y., CHERUBINI E. & GAIARSA J.L. A strychnine-insensitive modulation by glycine of GABA-mediated giant depolarizing potentials in immature rat hippocampal slices. (C) 112*P*

BEN ARI Y. & KRNJEVIĆ K. The ATP-sensitive K^+ channel opener diazoxide reduces anoxic depolarzation in rat CA3 hippocampal neurones in isolated slices. (PC) 192*P*

BENNETT K.A., LEMON R.N. & WERNER W. Indirect excitation of corticospinal neurones by intracortical stimulation in the conscious monkey. (C) 103*P*

BENNETT Max R. & LAVIDIS Nickolas A. The probability of quantal secretion at release sites in different calcium concentrations in toad (*Bufo marinus*) muscle. 219

BENNETT Max R., LAVIDIS Nickolas A. & LAVIDIS-ARMSON F. The probability of quantal secretion at release sites of different length in toad (*Bufo marinus*) muscle. 235

BENNETT P.B. Block of guinea-pig cardiac sodium channels by lidocaine and glycylxylidide. (PC) 191*P*

BENNETT T., GARDINER S.M. & TOMLINSON K.C. Cardiovascular and drinking responses to intracerebroventricular angiotensin II in streptozotocin-treated rats. (C) 82*P*

BERGFELD G. & FORRESTER T. Efflux of adenosine triphosphate from human erythrocytes in response to a brief pulse of hypoxia. (C) 88*P*

BERMAN N.J., DOUGLAS R.J. & MARTIN K.A.C. The conductances associated with inhibitory postsynaptic potentials are larger in visual cortical neurones *in vitro* than in similar neurones in intact, anaesthetized rats. (C) 107*P*

BERNENGO J.-C., LEMTIRI-CHLIEH F., OJEDA C. & PLATONOFF N. Measurement of cytosolic free Ca in cultured rat neonate and adult guinea-pig ventricular cardiac myocytes. (C) 45*P*

BERS Donald M. & HARRISON Simon M. Troponin-C replacement modifies the temperature-dependence of myofilament Ca-sensitivity of skinned cardiac muscle isolated from the rat. (PC) 166*P*

BERTHON B., CHAMPEIL P., CLARET M., COMBETTES L., DOUCET E. & ORLOWSKI S. Fast kinetics of calcium release induced by myo-inositoltrisphosphate in permeabilized rat hepatocytes. (PC) 178*P*

BETTANEY J., DRAY A. & FORSTER P. Resiniferatoxin: a capsaicin-like activator of peripheral fibres of the neonatal rat tail *in vitro* (PC) 150*P*

BETZ W.J., CHUA M. & RIDGE R.M.A.P. Inhibitory interactions between motoneurone terminals in neonatal rat lumbrical muscle. 25

BEVAN R.M. & BUTLER P.J. Oxygen consumption during voluntary diving in the tufted duck, *Aythya fuligula*, acclimated to summer and winter temperatures. (C) 132*P*

BIGELOW D.J., CLARKE D.M., INESI G., LOO T.W., MACLENNAN D.H. & SQUIER T.C. Correlation of function and molecular structure in the calcium transport ATPase of sarcoplasmic reticulum isolated from rabbit skeletal muscle. (C) 145*P*

BIGLAND-RITCHIE B., JOHANSSON R.S., THOMAS C.K. & WESTLING G. Twitch fusion in human motor units studied by motor axon stimulation. (C) 54*P*

BINDOKAS V.P. *see* ADAMS M.E., BINDOKAS V.P., DOLPHIN A.C. & SCOTT R.H. The inhibition of Ca^{2+} channel currents in cultured neonatal rat dorsal root ganglion (DRG) neurones by ω-Agatoxin-1A (a funnel web spider toxin). (C) 34*P*

BISCOE T.J. & DUCHEN M.R. Hypoxia raises intracellular calcium in Type I cells isolated from the rabbit carotid body. (C) 128*P*

BLACKSHAW Susanne E. & ROWLERSON Anthea M. ATPase activity and antibodies to vertebrate slow myosins distinguish fibre types in leech body wall muscle. (C) 51*P*

BLATTER L.A., BURI Arlette & MCGUIGAN J.A.S. Free intracellular magnesium concentration in isolated ferret ventricular muscle and in frog skeletal muscle measured with ion-selective microelectrodes containing the new magnesium sensor ETH 5214. (PC) 154*P*

BOISSONADE F.M. & MATTHEWS B. The effects of anaesthesia on the properties of neurones in the trigeminal brain-stem nuclei of the cat. (C) 115*P*

BOLTON T.B. *see* BEECH D.J. & BOLTON T.B. Two components of potassium current activated by depolarization of single smooth muscle cells from the rabbit portal vein. 293

BORGES R. *see* BALLESTA J.J., BORGES R., GARCÍA A.G. & HIDALGO M.J. Secretory and radioligand binding studies on muscarinic receptors in bovine and feline chromaffin cells. 411

BORN G.V.R. & SHAFI Shahida Evidence against the atherogenic passage of low-density lipoproteins between vascular endothelial cells in conscious rabbits. (C) 80*P*

BORON W.F., BOYARSKY G., GANZ M.B. & SCHLUE W.-R. The effect of DIDS on pH regulation of isolated single rat mesangial cells examined in the absence of HCO_3^-. (C) 125*P*

BOUNTRA C. & VAUGHAN-JONES R.D. Effect of intracellular and extracellular pH on contraction in isolated, mammalian cardiac muscle. 163

BOURNE G.B. & MCMAHON B.R. Control of cardiac output and its distribution in crustacean open circulatory systems. (C) 134*P*

BOWIE D. & SMART T.G. Properties of a kainate receptor expressed in *Xenopus laevis* oocytes after injection with bovine brain mRNA. (PC) 196*P*

BOYARSKY G. *see* BORON W.F., BOYARSKY G., GANZ M.B. & SCHLUE W.-R. The effect of DIDS on pH regulation of isolated single rat mesangial cells examined in the absence of HCO_3^-. (C) 125*P*

BRAUN M., HAYDON D.A. & MURRELL R.D. The effects of *n*-alcohols on the nicotinic acetylcholine receptor channel in cultured rat myotubes. (C) 37*P*

BRENNER B., FERENCZI M.A., IRVING M., KAPLAN J., SIMMONS R.M. & TOWNS-ANDREWS E. Structural changes upon activation of single isolated muscle fibres of the frog by photolysis of 'caged calcium' studied by time-resolved X-ray diffraction. (C) 59*P*

BRIDGE John H.B., SMOLLEY John R. & SPITZER Kenneth W. A discrete inward Na^+–Ca^{2+} exchange current accompanies the mechanical relaxation of isolated guinea-pig ventricular cells. (PC) 168*P*

BROWN E.R. *see* ABBOTT N. Joan & BROWN E.R. Effects of osmotic manipulation on extracellular routes for ion regulation in the squid Schwann cell. (C) 16*P*

BROWN J. & CZARNECKI A. Subtypes of specific binding of α-atrial natriuretic peptide (α-ANP) in rat brain. (C) 142*P*

BROWN M.C., LUNN E.R. & PERRY V.H. The route taken by axons growing down an intact distal nerve stump in mice. (PC) 148*P*

BROWN V., CARLISLE A.J. & MITCHELL M.A. Avian plasma concentrations of zinc; responses to stressors in the domestic fowl (*Gallus domesticus*). (C) 129*P*

BRYANT S.M., HART G. & RYDER Kathryn O. Captopril reduces L-type calcium current in isolated guinea-pig ventricular myocytes. (PC) 167*P*

BUCH J., COATS A., CONWAY J., MURPHY C. & SLEIGHT P. The baroreflex control of cardiac function in man. (C) 83*P*

BUCKINGHAM S.D., HARRISON J.B., LEECH C.A. & SATTELLE D.B. Nicotinic acetylcholine receptor activation in dissociated neurones from *Periplaneta americana* and *Musca domestica*. (PC) 193*P*

BULLIVANT E.M.A. & MUÑOZ D.J.B. A natriuretic substance is released from the kidney in response to low rates of saline infusion into the renal artery of the anaesthetized rat. (C) 141*P*

BURI Arlette *see* BLATTER L.A., BURI Arlette & MCGUIGAN J.A.S. Free intracellular magnesium concentration in isolated ferret ventricular muscle and in frog skeletal muscle measured with ion-selective microelectrodes containing the new magnesium sensor ETH 5214. (PC) 154*P*

BURKE E.P. & SANDERS K.M. Sodium-pump current in canine colonic myocytes is sufficient to provide a large contribution to membrane potential. (PC) 171*P*

BURKE W., EARLE J.W. & GOLDSMITH Robyn F. Transmitter metabolic changes in the decerebrate cat. (PC) 198*P*

BURTON K. The magnitude of the force rise after rapid shortening and restretch in fibres isolated from rabbit psoas muscle. (C) 66*P*

BUSTAMANTE J.C. *see* BARROS L.F., BUSTAMANTE J.C., JARVIS S.M. & YUDILEVICH D.L. Binding of

nitrobenzylthioinosine, an inhibitor of facilitated-diffusion nucleoside transport, to brush-border and basal membrane vesicles isolated from the human placenta. (PC) 179*P*

Butler P.J. *see* Anderson L.S., Butler P.J., Roberts C., Smale K., Snow D.H. & Woakes A.J. Partial pressures of arterial and end-tidal gases during graded exercise in thoroughbred race-horses. (C) 131*P*

Butler P.J. *see* Bevan R.M. & Butler P.J. Oxygen consumption during voluntary diving in the tufted duck, *Aythya fuligula*, acclimated to summer and winter temperatures. (C) 132*P*

Cachelin A.B. & Neuhaus R. The effect of Mg^{2+} on currents through single acetylcholine channels of rat PC12 cells. (C) 38*P*

Cady E.B., Jones D.A., Lynn J. & Newham D.J. Changes in force and intracellular metabolites during fatigue of human skeletal muscle. 311

Cady E.B., Elshove H., Jones D.A. & Moll A. The metabolic causes of slow relaxation in fatigued human skeletal muscle. 327

Camello P.J., Francis L.P., Madrid J.A., Pariente J.A., Salido G.M. & Singh J. Phorbol ester stimulates juice secretion and inhibits secretin-evoked secretory response in the pancreas of the anaesthetized rat. (C) 99*P*

Campbell E.A., Dray A., Perkins M.N. & Shaw W. Analgesic and anti-inflammatory activity of resiniferatoxin in the mouse and rat: comparison with capsaicin. (PC) 149*P*

Cannell M.B. *see* Adams D.J., van Breemen C., Cannell M.B. & Sage S.O. Na:Ca exchange in Fura-2-loaded bovine pulmonary artery endothelial cell monolayers. (PC) 183*P*

Carlisle A.J. *see* Brown V., Carlisle A.J. & Mitchell M.A. Avian plasma concentrations of zinc; responses to stressors in the domestic fowl (*Gallus domesticus*). (C) 129*P*

Case R.M., Seo Y. & Steward M.C. Regulation of intracellular pH in the isolated, perfused rabbit mandibular salivary gland studied by ^{31}P NMR spectroscopy. (C) 127*P*

Castellani L., Franzini-Armstrong C. & Loesser K. Shape, size and disposition of feet in junctions between transverse tubules and sarcoplasmic reticulum of Bivalvia, Insecta, Crustacea and Arachnida. (C) 118*P*

Cavieres J.D. & Walton T. Pre-steady-state kinetics of product release by Na,K-ATPase purified from pig kidney. (C) 90*P*

Cecchi G., Colomo F., Poggesi G. & Tesi C. A simple method to measure isometric twitch tension in single frog cardiac cells. (C) 49*P*

Cecchi G. *see* Ashley C.C., Bagni A., Cecchi G., Griffiths P.J. & Maeda Y. Time-resolved X-ray diffraction studies on intact single muscle fibres of *Rana temporaria*. (C) 58*P*

Cervero F. & Laird Jennifer M.A. Effects of reversible spinalization on the receptive field properties of dorsal horn neurones in anaesthetized rats. (C) 114*P*

Champeil P. *see* Berthon B., Champeil P., Claret M., Combettes L., Doucet E. & Orlowski S. Fast kinetics of calcium release induced by myo-inositoltrisphosphate in permeabilized rat hepatocytes. (PC) 178*P*

Chapman R.A. & Suleiman M.S. The effect of hypothermia on $[Na]_i$ and tension development during the Ca paradox in isolated ferret ventricular muscle. (C) 47*P*

Chernaeva L. The effect of neuropeptide Y on noradrenergic transmission in isolated guinea-pig vas deferens. (C) 76*P*

Cherubini E. *see* Ben Ari Y., Cherubini E. & Gaiarsa J.L. A strychnine-insensitive modulation by glycine of GABA-mediated giant depolarizing potentials in immature rat hippocampal slices. (C) 112*P*

Chodobski Adam & McKinley Michael J. Cerebral regulation of renal sodium excretion in sheep infused intravenously with hypertonic NaCl. 273

Christé G., De la Chapelle B., Lemtiri-Chlieh F. & Ojeda C. Presence of a Cs-resistant transient outward current which distorts I_{Ca} measurements in guinea-pig single ventricular cells. (PC) 169*P*

Chua M. *see* Betz W.J., Chua M. & Ridge R.M.A.P. Inhibitory interactions between motoneurone terminals in neonatal rat lumbrical muscle. 25

Claflin Dennis R., Faulkner John A. & Ward Christine E. Force–velocity relationships for isolated mouse soleus muscles during after-loaded and quick-released isotonic contractions. (PC) 161*P*

Clapp L.H. & Gurney A.M. Multiple effects of nitroprusside on membrane currents in smooth-muscle cells isolated from rabbit pulmonary artery. (C) 19*P*

Claret M. *see* Berthon B., Champeil P., Claret M., Combettes L., Doucet E. & Orlowski S. Fast kinetics of calcium release induced by myo-inositoltrisphosphate in permeabilized rat hepatocytes. (PC) 178*P*

CLARK H.W. & JACOBSON W. Detection of pulmonary surfactant in human amniotic fluid by polarized light microscopy. (C) 139P

CLARKE D.M. *see* BIGELOW D.J., CLARKE D.M., INESI G., LOO T.W., MACLENNAN D.H. & SQUIER T.C. Correlation of function and molecular structure in the calcium transport ATPase of sarcoplasmic reticulum isolated from rabbit skeletal muscle. (C) 145P

CLOWRY G.J. & MATTHEWS Margaret R. The effects of innervation of an inappropriate target, and exogenous ganglioside treatment, upon the retrograde axonal transport of HRP in axotomized hypoglossal neurones in the rat. (C) 73P

COATS A. *see* BUCH J., COATS A., CONWAY J., MURPHY C. & SLEIGHT P. The baroreflex control of cardiac function in man. (C) 83P

COHEN B.N., FAIN Gordon L. & FAIN Margery J. GABA and glycine channels in isolated ganglion cells from the goldfish retina 53

COHEN I. & TROMBA C. Inactivation of i_{k1} in isolated canine Purkinje myocytes is induced by isoprenaline but not blocked by propranolol. (C) 40P

COLAMARTINO G., MENINI A. & TORRE V. The blocking effect of divalent cations on the cyclic GMP-activated current in excised patches from retinal rods of the tiger salamander. (C) 122P

COLLMAN P.I. & DAVISON J.S. Vagal and splanchnic pathways are involved in the intestino-gastric inhibitory reflex in the anaesthetized ferret. (PC) 199P

COLOMO F. *see* CECCHI G., COLOMO F., POGGESI G. & TESI C. A simple method to measure isometric twitch tension in single frog cardiac cells. (C) 49P

COMBETTES L. *see* BERTHON B., CHAMPEIL P., CLARET M., COMBETTES L., DOUCET E. & ORLOWSKI S. Fast kinetics of calcium release induced by myo-inositoltrisphosphate in permeabilized rat hepatocytes. (PC) 178P

CONWAY J. *see* BUCH J., COATS A., CONWAY J., MURPHY C. & SLEIGHT P. The baroreflex control of cardiac function in man. (C) 83P

CULL-CANDY S.G., MATHIE A., SYMONDS C.J. & WYLLIE D.J.A. Distribution of quisqualate and kainate receptors in rat type-2 astrocytes and their progenitor cells in culture. (PC) 195P

CZARNECKI A. *see* BROWN J. & CZARNECKI A. Subtypes of specific binding of α-atrial natriuretic peptide (α-ANP) in rat brain. (C) 142P

DANTZIG J.A., GOLDMAN Y.E., LUTTMANN M.L., TRENTHAM D.R. & WOODWARD S.K.A. Binding of caged ATP diastereoisomers to rigor cross-bridges in glycerol-extracted fibres of rabbit psoas muscle. (C) 61P

DATTA A.K., HARRISON Linda M. & STEPHENS J.A. Task-dependent changes in the size of response to magnetic brain stimulation in human first dorsal interosseous muscle. 13

DAUNCEY M.J. & MOROVAT A. Cold-induced changes in nuclear 3,5,3′-triiodothyronine receptors of porcine skeletal muscle. (PC) 174P

DAUNCEY M.J., HOLDER G., INGRAM D.L., RUDD B.T. & SHAKESPEAR R.A. Thermal and nutritional regulation of insulin-like growth factor-I and cortisol in the young pig. (PC) 175P

DAUZENROTH M-E., DOLLY J.O., HABERMANN E., MOCHIDA S., POULAIN B., TAUC L., WADSWORTH J.D.F. & WELLER U. Light chain of tetanus toxin can be internalized into a cholinergic synapse of *Aplysia* by the heavy chain of botulinum neurotoxin and depresses the transmitter release. (C) 71P

DAVIES N.W., STANDEN N.B. & STANFIELD P.R. The kinetic behaviour of K_{ATP} channels in isolated membrane patches of frog skeletal muscle is regulated by internal pH. (PC) 188P

DAVISON J.S. *see* COLLMAN P.I. & DAVISON J.S. Vagal and splanchnic pathways are involved in the intestino-gastric inhibitory reflex in the anaesthetized ferret. (PC) 199P

DEARNLEY D.P., GOMEZ S., HALL M., LUSH D., POPE R. & WATSON M.S. A computerized system for analysing lung function data in man. (D) 4P

DE LA CHAPELLE B. *see* CHRISTÉ G., DE LA CHAPELLE B., LEMTIRI-CHLIEH F. & OJEDA C. Presence of a Cs-resistant transient outward current which distorts I_{Ca} measurements in guinea-pig single ventricular cells. (PC) 169P

DELAMERE Nicholas A. & PATERSON Christopher A. Studies on calcium transport in a rabbit lens membrane vesicle preparation. (PC) 181P

DOCHERTY R.J., FIEKERS J.F., WOOD J.N. & YEATS J.C. Histamine and prostaglandin D_2 modulate voltage-dependent calcium currents in adult rat sensory neurones in culture. (C) 36P

DOLLY J.O. *see* DAUZENROTH M-E., DOLLY J.O., HABERMANN E., MOCHIDA S., POULAIN B., TAUC L., WADSWORTH J.D.F. & WELLER U. Light chain of tetanus toxin can be internalized into a

cholinergic synapse of *Aplysia* by the heavy chain of botulinum neurotoxin and depresses the transmitter release. (C) 71*P*

DOLPHIN A.C. *see* ADAMS M.E., BINDOKAS V.P., DOLPHIN A.C. & SCOTT R.H. The inhibition of Ca^{2+} channel currents in cultured neonatal rat dorsal root ganglion (DRG) neurones by ω-Agatoxin-1A (a funnel web spider toxin). (C) 34*P*

DONOSO P., EISNER D.A. & O'NEILL S.C. Measurement of intracellular $[Na^+]$ in isolated rat cardiac myocytes using the fluorescent indicator SBFI. (C) 48*P*

DOUCET E. *see* BERTHON B., CHAMPEIL P., CLARET M., COMBETTES L., DOUCET E. & ORLOWSKI S. Fast kinetics of calcium release induced by myo-inositoltrisphosphate in permeabilized rat hepatocytes. (PC) 178*P*

DOUGLAS R.J. *see* BERMAN N.J. DOUGLAS R.J. & MARTIN K.A.C. The conductances associated with inhibitory postsynaptic potentials are larger in visual cortical neurones *in vitro* than in similar neurones in intact, anaesthetized rats. (C) 107*P*

DRAY A. *see* CAMPBELL E.A., DRAY A., PERKINS M.N. & SHAW W. Analgesic and anti-inflammatory activity of resiniferatoxin in the mouse and rat: comparison with capsaicin. (PC) 149*P*

DRAY A. *see* BETTANEY J., DRAY A. & FORSTER P. Resiniferatoxin: a capsaicin-like activator of peripheral fibres of the neonatal rat tail *in vitro* (PC) 150*P*

DUCHEN M.R. *see* BISCOE T.J. & DUCHEN M.R. Hypoxia raises intracellular calcium in Type I cells isolated from the rabbit carotid body. (C) 128*P*

DUGGAN K.A., HAWLEY C.M., MACDONALD G.J. & SHELLEY S. Sodium depletion decreases hepatic metabolism of vasoactive intestinal peptide in the rabbit 251

DUKES Iain D. & MORAD Martin Tedisamil increases the rate of inactivation of the transient outward current in single rat ventricular myocytes. (C) 33*P*

DUKES I.D. *see* AGUS Z.S., DUKES I.D. & MORAD M. Divalent cations modulate transient outward current in isolated rat ventricular myocytes. (C) 28*P*

DUNLOP W., GILLESPIE J.I., GREENWELL J.R. & OTUN H. Measurements of intracellular calcium (Ca_i^{2+}) and sodium (Na_i^+) in single cultured rat smooth muscle cells (A10) using the fluorescent dyes Fura-2 and SBFI. (PC) 201*P*

DUNNE M.J., GALLACHER D.V., PETERSEN O.H. & YULE D.I. The electrogenic sodium–alanine co-transport system in RINm5F insulin-secreting cells: patch-clamp and single cell $[Ca^{2+}]_i$ measurements. (C) 96*P*

EARLE J.W. *see* BURKE W., EARLE J.W. & GOLDSMITH Robyn F. Transmitter metabolic changes in the decerebrate cat. (PC) 198*P*

EARM Y.E., HO W.K. & SO I.S. Time course of Na–Ca exchange current in isolated rabbit atrial cells. (C) 41*P*

EDDY F.B., GRIERSON C., HAZON N. & SMITH N.F. Effects of atrial natriuretic peptide (ANP) on circulatory and ion regulatory physiology of rainbow trout (*Salmo gairdneri* Richardson) fed normal or high levels of dietary salt. (C) 137*P*

EDWARDS A.V. & JONES C.T. Adrenal responses to splanchnic nerve stimulation in conscious calves given naloxone. 339

EDWARDS R.H.T., GIBSON H. & GREGSON J.M. The effect of length on frequency-dependent force generation in human quadriceps muscle. (PC) 163*P*

EDWARDS R.H.T., GIBSON H. & GREGSON J.M. Length dependence of activation during isometric muscle fatigue in man. (PC) 164*P*

EISNER D.A. *see* DONOSO P., EISNER D.A. & O'NEILL S.C. Measurement of intracellular $[Na^+]$ in isolated rat cardiac myocytes using the fluorescent indicator SBFI. (C) 48*P*

EISNER D.A. *see* BARO I., EISNER D.A., FRY C.H., LISTON T.G., MONTGOMERY B.S.I. & RAIMBACH S.J. Measurement of intracellular pH in isolated ferret detrusor smooth muscle cells. (C) 124*P*

EKSTRÖM J., GARRETT J.R., MÅNSSON B. & PROCTOR G.B. Changes in protein composition of parotid saliva from anaesthetized rats after chronic sympathectomy, assessed by hydrophobic interaction chromatography. (C) 97*P*

ELSHOVE H. *see* CADY E.B., ELSHOVE H., JONES D.A. & MOLL A. The metabolic causes of slow relaxation in fatigued human skeletal muscle. 327

EVANS A.L., HARRISON Linda M. & STEPHENS J.A. Task-dependent changes in cutaneous reflexes recorded from various muscles controlling finger movement in man. 1

EYSTEINSSON Thor, HORIGUCHI Masayuki & KUNDAIKER Satish Spatial and temporal limitations of suppressive rod–cone interaction in *Xenopus* H-cells. (PC) 153*P*

FAIN Gordon L. *see* COHEN B.N., FAIN Gordon L. & FAIN Margery J. GABA and glycine channels in isolated ganglion cells from the goldfish retina 53

FAIN Gordon L. & FARAHBAKHSH Nasser A. Voltage-activated currents recorded from rabbit pigmented ciliary body epithelial cells in culture. 83

FAIN Margery J. *see* COHEN B.N., FAIN Gordon L. & FAIN Margery J. GABA and glycine channels in isolated ganglion cells from the goldfish retina 53

FARAHBAKHSH Nasser A. *see* FAIN Gordon L. & FARAHBAKHSH Nasser A. Voltage-activated currents recorded from rabbit pigmented ciliary body epithelial cells in culture. 83

FARMER S.F., INGRAM D.A., ROCHE S.W. & STEPHENS J.A. Evidence that mirror movements in the Klippel–Feil syndrome in man are produced by activity in branched-stem presynaptic inputs from corticospinal neurones. (C) 102*P*

FAULKNER John A. *see* CLAFLIN Dennis R., FAULKNER John A. & WARD Christine E. Force–velocity relationships for isolated mouse soleus muscles during after-loaded and quick-released isotonic contractions. (PC) 161*P*

FENTON M.J. & LEA T.J. Rapid activation of scallop skinned muscle fibres by laser photolysis of nitr-5. (C) 60*P*

FERENCZI M.A. *see* BRENNER B., FERENCZI M.A., IRVING M., KAPLAN J., SIMMONS R.M. & TOWNS-ANDREWS E. Structural changes upon activation of single isolated muscle fibres of the frog by photolysis of 'caged calcium' studied by time-resolved X-ray diffraction. (C) 59*P*

FERENCZI M.A., GOLDMAN Y.E. & TRENTHAM D.R. Relaxation of permeabilized, isolated muscle fibres of the rabbit by rapid chelation of Ca^{2+}-ions through laser pulse photolysis of 'caged-BAPTA'. (PC) 155*P*

FERN Robert & HARRISON P.J. The sequence of pressure-induced conduction failure of myelinated fibres in the isolated sciatic frog nerve (*Rana temporaria*) is dependent upon the pressure applied. (C) 77*P*

FERRELL William R. & SMITH Allan The effect of loading on position sense at the proximal interphalangeal joint of the human finger. 145

FIEKERS J.F. *see* DOCHERTY R.J., FIEKERS J.F., WOOD J.N. & YEATS J.C. Histamine and prostaglandin D_2 modulate voltage-dependent calcium currents in adult rat sensory neurones in culture. (C) 36*P*

FIELD M., GRENSPAN N.R., SULLIVAN S.K. & SWAMY K. Phorbol myristate acetate (PMA) inactivates an epithelial potassium channel from flounder intestine expressed by *Xenopus* oocytes. (C) 23*P*

FINK R.H.A. & HILLMAN G.M. Increased calcium release from the sarcoplasmic reticulum of normal and dystrophic isolated murine skeletal muscle in the presence of 4-aminopyridine. (PC) 156*P*

FLORES N.A., JUHASZ C., SHERIDAN D.J. & WINTERTON S.J. An IBM PC AT based system for the analysis of cardiac action potentials in real time. (D) 8*P*

FORRESTER T. *see* BERGFELD G. & FORRESTER T. Efflux of adenosine triphosphate from human erythrocytes in response to a brief pulse of hypoxia. (C) 88*P*

FORSTER I.C. & GREEFF N.G. The quantal gating charge of sodium-channel inactivation in the squid giant axon (*L. forbesii*). (C) 11*P*

FORSTER P. *see* BETTANEY J., DRAY A. & FORSTER P. Resiniferatoxin: a capsaicin-like activator of peripheral fibres of the neonatal rat tail *in vitro* (PC) 150*P*

FORTUNE N.S., GEEVES M.A. & RANATUNGA K.W. Tension transients initiated by pressure perturbation in isolated rabbit skinned muscle fibres. (PC) 158*P*

FRANCIS L.P. *see* CAMELLO P.J., FRANCIS L.P., MADRID J.A., PARIENTE J.A., SALIDO G.M. & SINGH J. Phorbol ester stimulates juice secretion and inhibits secretin-evoked secretory response in the pancreas of the anaesthetized rat. (C) 99*P*

FRANZINI-ARMSTRONG C. *see* CASTELLANI L., FRANZINI-ARMSTRONG C. & LOESSER K. Shape, size and disposition of feet in junctions between transverse tubules and sarcoplasmic reticulum of Bivalvia, Insecta, Crustacea and Arachnida. (C) 118*P*

FRY C.H. *see* BARO I., EISNER D.A., FRY C.H., LISTON T.G., MONTGOMERY B.S.I. & RAIMBACH S.J. Measurement of intracellular pH in isolated ferret detrusor smooth muscle cells. (C) 124*P*

GADSBY David C. *see* BAHINSKI Anthony, GADSBY David C., GREENGARD Paul & NAIRN Angus C. Chloride conductance regulated by protein kinase A in isolated guinea-pig ventricular myocytes. (C) 32*P*

GAGE Peter W., LAMB Graham D. & WAKEFIELD Bruce T. Transient and persistent sodium currents in normal and denervated mammalian skeletal muscle. 427

GAIARSA J.L. *see* BEN Ari Y., CHERUBINI E. & GAIARSA J.L. A strychnine-insensitive modulation by glycine of GABA-mediated giant depolarizing potentials in immature rat hippocampal slices. (C) 112*P*

GALLACHER D.V. *see* DUNNE M.J., GALLACHER D.V., PETERSEN O.H. & YULE D.I. The electrogenic sodium–alanine co-transport system in RINm5F insulin-secreting cells: patch-clamp and single cell $[Ca^{2+}]_i$ measurements. (C) 96*P*

GALLAGHER J., GARLICK P., HAMANN W., KEEVIL S., PETRICCIONE DI VADI P., PORTER D. & WEDLEY J.R. The use of magnetic resonance spectroscopy in assessing the efficacy of chemical lumbar sympathectomy in ischaemic vascular disease. (D) 3*P*

GALLAGHER Joel P. & HASUO Hiroshi Excitatory amino acid receptor-mediated EPSPs in rat dorsolateral septal nucleus neurones *in vitro*. 353

GALLAGHER Joel P. & HASUO Hiroshi Bicuculline- and phaclofen-sensitive components of *N*-methyl-D-aspartate-induced hyperpolarizations in rat dorsolateral septal nucleus neurones. 367

GAMBLE Theresa S., POPP David & STEWART Alex A.E. Time-resolved X-ray diffraction measurements of the isolated ABRM from *Mytilus edulis*. (PC) 204*P*

GANITKEVICH V. & ISEBERG G. Importance of current through L-type Ca-channels at the resting potential for the Ca balance of single coronary arterial myocytes of the guinea-pig. (C) 30*P*

GANZ M.B. *see* BORON W.F., BOYARSKY G., GANZ M.B. & SCHLUE W.-R. The effect of DIDS on pH regulation of isolated single rat mesangial cells examined in the absence of HCO_3^-. (C) 125*P*

GARCÍA A.G. *see* BALLESTA J.J., BORGES R., GARCÍA A.G. & HIDALGO M.J. Secretory and radioligand binding studies on muscarinic receptors in bovine and feline chromaffin cells. 411

GARDINER S.M. *see* BENNETT T., GARDINER S.M. & TOMLINSON K.C. Cardiovascular and drinking responses to intracerebroventricular angiotensin II in streptozotocin-treated rats. (C) 82*P*

GARDNER-MEDWIN A.R. *see* BARKLEY G.L., GARDNER-MEDWIN A.R., MORAN J., NAGEL-LEIBY S., SIMKINS R.T., TEPLEY N. & WELCH K.M.A. Magnetic fields during spreading depression in the cortex of anaesthetized rabbits. (C) 101*P*

GARLAND C.J. Influence of the endothelium and α-adrenoreceptor antagonists on responses to noradrenaline in the rabbit basilar artery 205

GARLICK P. *see* GALLAGHER J., GARLICK P., HAMANN W., KEEVIL S., PETRICCIONE DI VADI P., PORTER D. & WEDLEY J.R. The use of magnetic resonance spectroscopy in assessing the efficacy of chemical lumbar sympathectomy in ischaemic vascular disease. (D) 3*P*

GARRETT J.R. *see* EKSTRÖM J., GARRETT J.R., MÅNSSON B. & PROCTOR G.B. Changes in protein composition of parotid saliva from anaesthetized rats after chronic sympathectomy, assessed by hydrophobic interaction chromatography. (C) 97*P*

GASSER R. & VAUGHAN-JONES R.D. Myocardial ischaemia: release of potassium can be inhibited by tolbutamide and glibenclamide in isolated guinea-pig papillary muscle. (C) 43*P*

GEE H. & TAYLOR E.W. Detection of sinus arrhythmia in the human fetus *in utero* using ultrasound and Doppler blood-flow waveforms in the major vessels. (PC) 184*P*

GEEVES M.A. *see* FORTUNE N.S., GEEVES M.A. & RANATUNGA K.W. Tension transients initiated by pressure perturbation in isolated rabbit skinned muscle fibres. (PC) 158*P*

GIBSON H. *see* EDWARDS R.H.T., GIBSON H. & GREGSON J.M. The effect of length on frequency-dependent force generation in human quadriceps muscle. (PC) 163*P*

GIBSON H. *see* EDWARDS R.H.T., GIBSON H. & GREGSON J.M. Length dependence of activation during isometric muscle fatigue in man. (PC) 164*P*

GILLESPIE J.I. *see* DUNLOP W., GILLESPIE J.I., GREENWELL J.R. & OTUN H. Measurements of intracellular calcium (Ca_i^{2+}) and sodium (Na_i^+) in single cultured rat smooth muscle cells (A10) using the fluorescent dyes Fura-2 and SBFI. (PC) 201*P*

GIRALDEZ F., SEPÚLVEDA F.V., SHEPPARD D.N. & VALVERDE M.A. Potassium currents in isolated *Necturus* enterocytes. (C) 13*P*

GODT R.E. & KENTISH J.C. Effect of pH on isometric force, MgATPase and tension-cost of skinned skeletal muscle fibres of the rabbit and cardiac muscles from rat. (C) 68*P*

GOLDMAN Y.E. *see* DANTZIG J.A., GOLDMAN Y.E., LUTTMANN M.L., TRENTHAM D.R. & WOODWARD S.K.A. Binding of caged ATP diastereoisomers to rigor cross-bridges in glycerol-extracted fibres of rabbit psoas muscle. (C) 61*P*

GOLDMAN Y.E. *see* FERENCZI M.A., GOLDMAN Y.E. & TRENTHAM D.R. Relaxation of permeabilized, isolated muscle fibres of the rabbit by rapid chelation of Ca^{2+}-ions through laser pulse photolysis of 'caged-BAPTA'. (PC) 155*P*

GOLDSMITH Robyn F. *see* BURKE W., EARLE J.W. & GOLDSMITH Robyn F. Transmitter metabolic changes in the decerebrate cat. (PC) 198*P*

GOLDSPINK Geoffrey & WILLIAMS Pamela E. The effect of immobilization on gene expression and post-natal determination of muscle fibre type in the mouse soleus muscle. (C) 52*P*

GOMEZ K. & TAYLOR A. Spectral analysis of the demodulated EMG in isometric and isotonic human muscle contractions. (C) 105*P*

GOMEZ S. *see* DEARNLEY D.P., GOMEZ S., HALL M., LUSH D., POPE R. & WATSON M.S. A computerized system for analysing lung function data in man. (D) 4*P*

GONG Q.L. & LI P. Inhibitory effect of nucleus raphe obscurus on the pressor response to stimulation of the hypothalamic and midbrain defence areas in the anaesthetized rabbit. (C) 85*P*

GONZALEZ-SERRATOS H., INESI G. & ORTEGA Alicia Effects of D-600 on isolated fibres and isolated sarcoplasmic reticulum of frog skeletal muscle. (C) 119*P*

GORDON A.M. *see* ALLEN T.StC. & GORDON A.M. Cycling cross-bridges produce structural changes in Troponin-C in demembranated fibres of rabbit psoas muscle. (C) 63*P*

GREEFF N.G. *see* FORSTER I.C. & GREEFF N.G. The quantal gating charge of sodium-channel inactivation in the squid giant axon (*L. forbesii*). (C) 11*P*

GREENGARD Paul *see* BAHINSKI Anthony, GADSBY David C., GREENGARD Paul & NAIRN Angus C. Chloride conductance regulated by protein kinase A in isolated guinea-pig ventricular myocytes. (C) 32*P*

GREENWELL J.R. *see* DUNLOP W., GILLESPIE J.I., GREENWELL J.R. & OTUN H. Measurements of intracellular calcium (Ca_i^{2+}) and sodium (Na_i^{+}) in single cultured rat smooth muscle cells (A10) using the fluorescent dyes Fura-2 and SBFI. (PC) 201*P*

GREGSON J.M. *see* EDWARDS R.H.T., GIBSON H. & GREGSON J.M. The effect of length on frequency-dependent force generation in human quadriceps muscle. (PC) 163*P*

GREGSON J.M. *see* EDWARDS R.H.T., GIBSON H. & GREGSON J.M. Length dependence of activation during isometric muscle fatigue in man. (PC) 164*P*

GRENSPAN N.R. *see* FIELD M., GRENSPAN N.R., SULLIVAN S.K. & SWAMY K. Phorbol myristate acetate (PMA) inactivates an epithelial potassium channel from flounder intestine expressed by *Xenopus* oocytes. (C) 23*P*

GRIERSON C. *see* EDDY F.B., GRIERSON C., HAZON N. & SMITH N.F. Effects of atrial natriuretic peptide (ANP) on circulatory and ion regulatory physiology of rainbow trout (*Salmo gairdneri* Richardson) fed normal or high levels of dietary salt. (C) 137*P*

GRIFFITHS P.J. *see* ASHLEY C.C., BAGNI A., CECCHI G., GRIFFITHS P.J. & MAEDA Y. Time-resolved X-ray diffraction studies on intact single muscle fibres of *Rana temporaria*. (C) 58*P*

GRINNELL Alan D. & PAWSON Peter A. Dependence of spontaneous release at frog junctions on synaptic strength, external calcium and terminal length. 397

GUNASENA Karen T., MORRISON J.F.B., NIMMO A.J. & WHITAKER Elaine M. The effect of oestradiol and progesterone on neurokinin receptors in rat uterus. (C) 75*P*

GURNEY A.M. *see* CLAPP L.H. & GURNEY A.M. Multiple effects of nitroprusside on membrane currents in smooth-muscle cells isolated from rabbit pulmonary artery. (C) 19*P*

HABERMANN E. *see* DAUZENROTH M-E., DOLLY J.O., HABERMANN E., MOCHIDA S., POULAIN B., TAUC L., WADSWORTH J.D.F. & WELLER U. Light chain of tetanus toxin can be internalized into a cholinergic synapse of *Aplysia* by the heavy chain of botulinum neurotoxin and depresses the transmitter release. (C) 71*P*

HADLEY R.W. & LEDERER W.J. L-type calcium channel gating current in isolated guinea-pig ventricular myocytes. (C) 27*P*

HALL A.C., HARVEY C.M., HENDRY B.M. & LUCIO F.J. Effects of hydrostatic pressure and membrane cholesterol depletion on L-lysine transport in human erythrocytes. (C) 89*P*

HALL M. *see* DEARNLEY D.P., GOMEZ S., HALL M., LUSH D., POPE R. & WATSON M.S. A computerized system for analysing lung function data in man. (D) 4*P*

HAMANN W. *see* GALLAGHER J., GARLICK P., HAMANN W., KEEVIL S., PETRICCIONE DI VADI P., PORTER D. & WEDLEY J.R. The use of magnetic resonance spectroscopy in assessing the efficacy of chemical lumbar sympathectomy in ischaemic vascular disease. (D) 3*P*

HANKINS M.W. & IKEDA H. Properties of excitatory amino acid receptors on rod driven horizontal cells in the isolated rat retina. (C) 109*P*

HARDIE R.C. Ionic channels in dissociated retinal neurones of the blowfly. (PC) 203*P*

HARPER A.A. *see* BACKHOUSE S.S., HARPER A.A., VAN VLIET B.N. & WEST N.H. Localization and conduction velocities of arterial baroreceptors in the anaesthetized snake. (C) 135*P*

HARRIS H.W. Jr, JANOSHAZI A., KIKERI D., SOLOMON A.K. & ZEIDEL M.L. Proton permeability of toad bladder aggrephore membranes: role of the water channel in proton transport. (C) 145*P*

HARRISON J.B. *see* BUCKINGHAM S.D., HARRISON J.B., LEECH C.A. & SATTELLE D.B. Nicotinic acetylcholine receptor activation in dissociated neurones from *Periplaneta americana* and *Musca domestica*. (PC) 193*P*

HARRISON Linda M. *see* EVANS A.L., HARRISON Linda M. & STEPHENS J.A. Task-dependent changes in cutaneous reflexes recorded from various muscles controlling finger movement in man. 1

HARRISON Linda M. *see* DATTA A.K., HARRISON Linda M. & STEPHENS J.A. Task-dependent changes in the size of response to magnetic brain stimulation in human first dorsal interosseous muscle. 13

HARRISON Linda M., IRONTON R. & STEPHENS J.A. Cross correlation analysis of multi-unit EMG recordings in man. (D) 6*P*

HARRISON P.J. *see* FERN Robert & HARRISON P.J. The sequence of pressure-induced conduction failure of myelinated fibres in the isolated sciatic frog nerve (*Rana temporaria*) is dependent upon the pressure applied. (C) 77*P*

HARRISON Simon M. *see* BERS Donald M. & HARRISON Simon M. Troponin-C replacement modifies the temperature-dependence of myofilament Ca-sensitivity of skinned cardiac muscle isolated from the rat. (PC) 166*P*

HART G. *see* BRYANT S.M., HART G. & RYDER Kathryn O. Captopril reduces L-type calcium current in isolated guinea-pig ventricular myocytes. (PC) 167*P*

HARVEY Brian J. Water fluxes and conductive proton passage across apical and basolateral cell membranes of oxytocin-treated isolated frog skin and urinary bladder: evidence for a common pathway. (C) 91*P*

HARVEY C.M. *see* HALL A.C., HARVEY C.M., HENDRY B.M. & LUCIO F.J. Effects of hydrostatic pressure and membrane cholesterol depletion on L-lysine transport in human erythrocytes. (C) 89*P*

HARVEY R.D. & HUME J.R. Isoprenaline activates a chloride conductance in isolated guinea-pig and rabbit ventricular myocytes. (C) 39*P*

HASUO Hiroshi *see* GALLAGHER Joel P. & HASUO Hiroshi Excitatory amino acid receptor-mediated EPSPs in rat dorsolateral septal nucleus neurones *in vitro*. 353

HASUO Hiroshi *see* GALLAGHER Joel P. & HASUO Hiroshi Bicuculline- and phaclofen-sensitive components of *N*-methyl-D-aspartate-induced hyperpolarizations in rat dorsolateral septal nucleus neurones. 367

HAWLEY C.M. *see* DUGGAN K.A., HAWLEY C.M., MACDONALD G.J. & SHELLEY S. Sodium depletion decreases hepatic metabolism of vasoactive intestinal peptide in the rabbit 251

HAYDON D.A. *see* BRAUN M., HAYDON D.A. & MURRELL R.D. The effects of *n*-alcohols on the nicotinic acetylcholine receptor channel in cultured rat myotubes. (C) 37*P*

HAYDON P.G., LUKOWIAK K., MAN-SON-HING H.J. & ZORAN M.J. Mechanisms of synaptic plasticity: modulation of secretory apparatus in cultured neuronal somata of *Helisoma trivolvis*. (C) 111*P*

HAYDON P.G., LUKOWIAK K., MAN-SON-HING H.J. & ZORAN M.J. Mechanisms of synaptic plasticity: Use of a giant soma–soma synapse in cultured neuronal somata of *Helisoma trivolvis*. (PC) 147*P*

HAZON N. *see* EDDY F.B., GRIERSON C., HAZON N. & SMITH N.F. Effects of atrial natriuretic peptide (ANP) on circulatory and ion regulatory physiology of rainbow trout (*Salmo gairdneri* Richardson) fed normal or high levels of dietary salt. (C) 137*P*

HENDRY B.M. *see* HALL A.C., HARVEY C.M., HENDRY B.M. & LUCIO F.J. Effects of hydrostatic pressure and membrane cholesterol depletion on L-lysine transport in human erythrocytes. (C) 89*P*

HIDALGO M.J. *see* BALLESTA J.J., BORGES R., GARCÍA A.G. & HIDALGO M.J. Secretory and radioligand binding studies on muscarinic receptors in bovine and feline chromaffin cells. 411

HILLMAN G.M. *see* FINK R.H.A. & HILLMAN G.M. Increased calcium release from the sarcoplasmic reticulum of normal and dystrophic isolated murine skeletal muscle in the presence of 4-aminopyridine. (PC) 156*P*

HO W.K. *see* EARM Y.E., HO W.K. & SO I.S. Time course of Na–Ca exchange current in isolated rabbit atrial cells. (C) 41*P*

HOLDEN-DYE L., PARRI H.R. & WALKER R.J. Anion selectivity of the GABA-operated channel in somatic muscle cells of the parasitic nematode *Ascaris suum*. (PC) 194*P*

HOLDER G. *see* DAUNCEY M.J., HOLDER G., INGRAM D.L., RUDD B.T. & SHAKESPEAR R.A. Thermal and nutritional regulation of insulin-like growth factor-I and cortisol in the young pig. (PC) 175*P*

HOLLINGWORTH S. *see* BAYLOR S.M., HOLLINGWORTH S. & KONISHI M. Calcium transients in intact single frog skeletal muscle fibres measured with the fluorescence indicator dye Mag-fura-2. (C) 69*P*

HOLMBERG S.R.M. & WILLIAMS A.J. Effects of doxorubicin and mitoxantrone on single calcium release channels from isolated sheep cardiac sarcoplasmic reticulum. (PC) 202*P*

HOMSHER E. & MILLAR N.C. Tension transients induced by photolysis of caged phosphate in glycerinated rabbit soleus muscle fibres. (C) 62*P*

HORIGUCHI Masayuki *see* EYSTEINSSON Thor, HORIGUCHI Masayuki & KUNDAIKER Satish Spatial and temporal limitations of suppressive rod–cone interaction in *Xenopus* H-cells. (PC) 153*P*

HORN N.M. *see* AIKEN S.P., HORN N.M. & SAUNDERS N.R. Effect of histidine on zinc distribution in anaesthetized rats. (C) 144*P*

HOU T.-t., JOHNSON J.D. & RALL J.A. Mg^{2+} and Ca^{2+} dissociation rates from frog parvalbumin at 0 °C: implications for role of parvalbumin in skeletal muscle relaxation. (C) 120*P*

HUANG C.L.-H. & PEACHEY L.D. Charge movements during a frog skeletal muscle action potential. (C) 121*P*

HUME J.R. *see* HARVEY R.D. & HUME J.R. Isoprenaline activates a chloride conductance in isolated guinea-pig and rabbit ventricular myocytes. (C) 39*P*

HUME J.R. & LEBLANC N. Sarcoplasmic reticulum Ca^{2+} release in guinea-pig ventricular myocytes in the absence of Ca^{2+} current. (C) 44*P*

HUNTER M. & HURST A.M. Acid-inhibition of channels in the apical membrane of isolated frog diluting segment. (PC) 180*P*

HURST A.M. *see* HUNTER M. & HURST A.M. Acid-inhibition of channels in the apical membrane of isolated frog diluting segment. (PC) 180*P*

HUXLEY Virginia H. & MEYER D.J. Jr Volume status dependence of atrial peptide-induced changes in capillary water conductance: evidence from individually perfused mesenteric microvessels of decerebrate leopard frogs. (C) 78*P*

IKEDA H. *see* HANKINS M.W. & IKEDA H. Properties of excitatory amino acid receptors on rod driven horizontal cells in the isolated rat retina. (C) 109*P*

INESI G. *see* GONZALEZ-SERRATOS H., INESI G. & ORTEGA Alicia Effects of D-600 on isolated fibres and isolated sarcoplasmic reticulum of frog skeletal muscle. (C) 119*P*

INESI G. *see* BIGELOW D.J., CLARKE D.M., INESI G., LOO T.W., MACLENNAN D.H. & SQUIER T.C. Correlation of function and molecular structure in the calcium transport ATPase of sarcoplasmic reticulum isolated from rabbit skeletal muscle. (C) 145*P*

INGRAM D.A. *see* FARMER S.F., INGRAM D.A., ROCHE S.W. & STEPHENS J.A. Evidence that mirror movements in the Klippel–Feil syndrome in man are produced by activity in branched-stem presynaptic inputs from corticospinal neurones. (C) 102*P*

INGRAM D.L. *see* DAUNCEY M.J., HOLDER G., INGRAM D.L., RUDD B.T. & SHAKESPEAR R.A. Thermal and nutritional regulation of insulin-like growth factor-I and cortisol in the young pig. (PC) 175*P*

INOUÉ Shinya & OLDENBOURG Rudolf Polarized light microscopy of striated muscle and analysis of edge birefringence. (C) 56*P*

IRONTON R. *see* HARRISON Linda M., IRONTON R. & STEPHENS J.A. Cross correlation analysis of multi-unit EMG recordings in man. (D) 6*P*

IRVING M., KENDRICK-JONES J., SHRIMPTON C. & SLEEP J. The orientation of rhodamine probes attached to myosin light chain-2 (LC-2) in relaxing and rigor conditions in single fibres isolated from rabbit psoas muscle. (C) 57*P*

IRVING M. *see* BRENNER B., FERENCZI M.A., IRVING M., KAPLAN J., SIMMONS R.M. & TOWNS-ANDREWS E. Structural changes upon activation of single isolated muscle fibres of the frog by photolysis of 'caged calcium' studied by time-resolved X-ray diffraction. (C) 59*P*

ISEBERG G. *see* GANITKEVICH V. & ISEBERG G. Importance of current through L-type Ca-channels at the resting potential for the Ca balance of single coronary arterial myocytes of the guinea-pig. (C) 30*P*

IWATA A. *see* ALMERS W., IWATA A., SPRUCE A.E. & WHITE J.M. Patch clamp studies of fusion of human erythrocytes to murine fibroblasts mediated by influenza virus fusion protein. (C) 95*P*

JACKSON A.A., SLEVIN K. & WATERLOW J.C. A model of whole body protein turnover in man that includes a pool turning over by life-time kinetics. (C) 143*P*

JACOBSON W. *see* CLARK H.W. & JACOBSON W. Detection of pulmonary surfactant in human amniotic fluid by polarized light microscopy. (C) 139*P*

JAHR C.E. & LESTER R.A.J. Depression of calcium currents by quisqualic acid in neonatal rat hippocampal neurones in culture. (PC) 197*P*

JANOSHAZI A. *see* HARRIS H.W. Jr, JANOSHAZI A., KIKERI D., SOLOMON A.K. & ZEIDEL M.L. Proton permeability of toad bladder aggrephore membranes: role of the water channel in proton transport. (C) 145*P*

JARVIS S.M. *see* BARROS L.F., BUSTAMANTE J.C., JARVIS S.M. & YUDILEVICH D.L. Binding of nitrobenzylthioinosine, an inhibitor of facilitated-diffusion nucleoside transport, to brush-border and basal membrane vesicles isolated from the human placenta. (PC) 179*P*

JOHANSSON R.S. *see* BIGLAND-RITCHIE B., JOHANSSON R.S., THOMAS C.K. & WESTLING G. Twitch fusion in human motor units studied by motor axon stimulation. (C) 54*P*

JOHNSON J.D. *see* HOU T.-t., JOHNSON J.D. & RALL J.A. Mg^{2+} and Ca^{2+} dissociation rates from frog parvalbumin at 0 °C: implications for role of parvalbumin in skeletal muscle relaxation. (C) 120*P*

JOHNSON M.S., MACEWAN D.J., MITCHELL R. & THOMSON F.J. Lithium enhances the inhibitory effect of protein kinase C on LHRH-induced $^{45}Ca^{2+}$ influx into rat anterior pituitary tissue *in vitro*. (PC) 185*P*

JOHNSON M.S., MACEWAN D.J. & MITCHELL R. Depolarization-induced $^{45}Ca^{2+}$ influx into rat anterior pituitary cells *in vitro* can be modulated in two ways by protein kinase C. (PC) 186*P*

JONES C.T. *see* EDWARDS A.V. & JONES C.T. Adrenal responses to splanchnic nerve stimulation in conscious calves given naloxone. 339

JONES D.A. *see* CADY E.B., JONES D.A., LYNN J. & NEWHAM D.J. Changes in force and intracellular metabolites during fatigue of human skeletal muscle. 311

JONES D.A. *see* CADY E.B., ELSHOVE H., JONES D.A. & MOLL A. The metabolic causes of slow relaxation in fatigued human skeletal muscle. 327

JUHASZ C. *see* FLORES N.A., JUHASZ C., SHERIDAN D.J. & WINTERTON S.J. An IBM PC AT based system for the analysis of cardiac action potentials in real time. (D) 8*P*

KAO C.Y., WAKUI M., WANG S.Y. & YOSHINO M. The outward current of the isolated rat myometrium. (C) 20*P*

KAPLAN J. *see* BRENNER B., FERENCZI M.A., IRVING M., KAPLAN J., SIMMONS R.M. & TOWNS-ANDREWS E. Structural changes upon activation of single isolated muscle fibres of the frog by photolysis of 'caged calcium' studied by time-resolved X-ray diffraction. (C) 59*P*

KARSCHIN Andreas & LIPTON Stuart A. Calcium channels in solitary retinal ganglion cells from post-natal rat. 379

KAWAI M. & MARCUSSEN B. Kinetics of MgATP binding, oscillatory power production, and the rate-limiting step in isolated insect muscle fibres. (C) 64*P*

KEEVIL S. *see* GALLAGHER J., GARLICK P., HAMANN W., KEEVIL S., PETRICCIONE DI VADI P., PORTER D. & WEDLEY J.R. The use of magnetic resonance spectroscopy in assessing the efficacy of chemical lumbar sympathectomy in ischaemic vascular disease. (D) 3*P*

KELLY J.S. & PENINGTON N.J. 5-Hydroxytryptamine inhibits the voltage-dependent calcium current of acutely dissociated central neurones from the adult rat dorsal raphe nucleus. (C) 35*P*

KENDRICK-JONES J. *see* IRVING M., KENDRICK-JONES J., SHRIMPTON C. & SLEEP J. The orientation of rhodamine probes attached to myosin light chain-2 (LC-2) in relaxing and rigor conditions in single fibres isolated from rabbit psoas muscle. (C) 57*P*

KENTISH J.C. *see* GODT R.E. & KENTISH J.C. Effect of pH on isometric force, MgATPase and tension-cost of skinned skeletal muscle fibres of the rabbit and cardiac muscles from rat. (C) 68*P*

KEYNES R.D. An 8-state 5-stage model of the voltage-gated sodium channel. (C) 12*P*

KIKERI D. *see* HARRIS H.W. Jr, JANOSHAZI A., KIKERI D., SOLOMON A.K. & ZEIDEL M.L. Proton permeability of toad bladder aggrephore membranes: role of the water channel in proton transport. (C) 145*P*

KIRCHHOFF C.G., REEH P.W. & WADDELL P.J. Sensory endings of C- and A-fibres are differentially sensitive to tetrodotoxin in the rat skin *in vitro*. (C) 116*P*

KIRKWOOD P.A. & SEARS T.A. Field potentials in the contralateral ventral horn evoked by individual thoracic respiratory interneurones in the anaesthetized cat. (C) 106*P*

KONISHI M. *see* BAYLOR S.M., HOLLINGWORTH S. & KONISHI M. Calcium transients in intact single frog skeletal muscle fibres measured with the fluorescence indicator dye Mag-fura-2. (C) 69*P*

KRNJEVIĆ K. *see* BEN ARI Y. & KRNJEVIĆ K. The ATP-sensitive K^+ channel opener diazoxide reduces anoxic depolarzation in rat CA3 hippocampal neurones in isolated slices. (PC) 192*P*

KROESE A.B.A., PRINS M. & SCHELLART N.A.M. Regional differences in conduction velocity and fibre diameter in posterior lateral line afferent axons in the trout. (C) 136*P*

KUNDAIKER Satish *see* EYSTEINSSON Thor, HORIGUCHI Masayuki & KUNDAIKER Satish Spatial and temporal limitations of suppressive rod–cone interaction in *Xenopus* H-cells. (PC) 153*P*

LAIRD Jennifer M.A. *see* CERVERO F. & LAIRD Jennifer M.A. Effects of reversible spinalization on the receptive field properties of dorsal horn neurones in anaesthetized rats. (C) 114*P*

LAMB G.D. & STEPHENSON D.G. Activation and reduction of calcium release in skinned muscle fibres from the cane toad. (C) 70*P*

LAMB Graham D. *see* GAGE Peter W., LAMB Graham D. & WAKEFIELD Bruce T. Transient and persistent sodium currents in normal and denervated mammalian skeletal muscle. 427

LAUGHLIN S.B. & WECKSTRÖM M. The activation of a slow voltage-depedent potassium conductance is crucial for light adaptation in blowfly receptos. (PC) 200*P*

LAVIDIS Nickolas A. *see* BENNETT Max R. & LAVIDIS Nickolas A. The probability of quantal secretion at release sites in different calcium concentrations in toad (*Bufo marinus*) muscle. 219

LAVIDIS Nickolas A. *see* BENNETT Max R., LAVIDIS Nickolas A. & LAVIDIS-ARMSON F. The probability of quantal secretion at release sites of different length in toad (*Bufo marinus*) muscle. 235

LAVIDIS-ARMSON F. *see* BENNETT Max R., LAVIDIS Nickolas A. & LAVIDIS-ARMSON F. The probability of quantal secretion at release sites of different length in toad (*Bufo marinus*) muscle. 235

LEA T.J. *see* FENTON M.J. & LEA T.J. Rapid activation of scallop skinned muscle fibres by laser photolysis of nitr-5. (C) 60*P*

LEBLANC N. *see* HUME J.R. & LEBLANC N. Sarcroplasmic reticulum Ca^{2+} release in guinea-pig ventricular myocytes in the absence of Ca^{2+} current. (C) 44*P*

LEDERER W.J. *see* HADLEY R.W. & LEDERER W.J. L-type calcium channel gating current in isolated guinea-pig ventricular myocytes. (C) 27*P*

LEE K.S. Potentiation of the L-type Ca^{2+} channels is critically dependent on the resting membrane potentials in single guinea-pig ventricular cells. (C) 29*P*

LEE K.S. & TSAI T.D. Open channel block of the fast inward sodium current by TTX in single monkey heart cells ? (PC) 190*P*

LEECH C.A. *see* BUCKINGHAM S.D., HARRISON J.B., LEECH C.A. & SATTELLE D.B. Nicotinic acetylcholine receptor activation in dissociated neurones from *Periplaneta americana* and *Musca domestica*. (PC) 193*P*

LEMON R.N. *see* BENNETT K.A., LEMON R.N. & WERNER W. Indirect excitation of corticospinal neurones by intracortical stimulation in the conscious monkey. (C) 103*P*

LEMTIRI-CHLIEH F. *see* BERNENGO J.-C., LEMTIRI-CHLIEH F., OJEDA C. & PLATONOFF N. Measurement of cytosolic free Ca in cultured rat neonate and adult guinea-pig ventricular cardiac myocytes. (C) 45*P*

LEMTIRI-CHLIEH F. *see* CHRISTÉ G., DE LA CHAPELLE B., LEMTIRI-CHLIEH F. & OJEDA C. Presence of a Cs-resistant transient outward current which distorts I_{Ca} measurements in guinea-pig single ventricular cells. (PC) 169*P*

LEOTY C. & NOIREAUD J. Two components of potassium contractures in isolated ferret cremaster fibres. (PC) 172*P*

LESTER R.A.J. *see* JAHR C.E. & LESTER R.A.J. Depression of calcium currents by quisqualic acid in neonatal rat hippocampal neurones in culture. (PC) 197*P*

LEUCHTAG H. Richard Na-channel gating as a ferroelectric-superionic phase transition. (C) 10*P*

LEVI Allan J. A reduction of calcium current partly underlies action potential shortening with strophanthidin in isolated ventricular myocytes from the guinea-pig. (C) 42*P*

LEVINE Rhea J.C. Relaxed thick filaments from vertebrate and invertebrate striated muscles have overlapping myosin heads. (PC) 157*P*

LEVY A. & LIGHTMAN Stafford L. Posterior pituitary vasopressin mRNA in salt loaded rats: effects of pituitary stalk transection. (C) 140*P*

LI P. *see* GONG Q.L. & LI P. Inhibitory effect of nucleus raphe obscurus on the pressor response to stimulation of the hypothalamic and midbrain defence areas in the anaesthetized rabbit. (C) 85*P*

LIGHTMAN Stafford L. *see* LEVY A. & LIGHTMAN Stafford L. Posterior pituitary vasopressin mRNA in salt loaded rats: effects of pituitary stalk transection. (C) 140*P*

LIPTON Stuart A. *see* KARSCHIN Andreas & LIPTON Stuart A. Calcium channels in solitary retinal ganglion cells from post-natal rat. 379

LISTON T.G. *see* BARO I., EISNER D.A., FRY C.H., LISTON T.G., MONTGOMERY B.S.I. & RAIMBACH S.J. Measurement of intracellular pH in isolated ferret detrusor smooth muscle cells. (C) 124*P*

LLEWELLYN M., PROCHAZKA A. & YANG J.F. Human H-reflexes are reduced in difficult beam-walking compared to stance and normal walking. (C) 104*P*

LLEWELLYN M., PROCHAZKA A. & TREND P. Control of ankle extensor activation prior to and after foot contact in cat step cycle. (PC) 146*P*

LOESSER K. *see* CASTELLANI L., FRANZINI-ARMSTRONG C. & LOESSER K. Shape, size and disposition of feet in junctions between transverse tubules and sarcoplasmic reticulum of Bivalvia, Insecta, Crustacea and Arachnida. (C) 118*P*

LOMBARDI V. & PIAZZESI G. Mechanical properties of tetanized frog muscle fibres during the transitory phase of force response to lengthening. (C) 65*P*

LOO T.W. *see* BIGELOW D.J., CLARKE D.M., INESI G., LOO T.W., MACLENNAN D.H. & SQUIER T.C. Correlation of function and molecular structure in the calcium transport ATPase of sarcoplasmic reticulum isolated from rabbit skeletal muscle. (C) 145*P*

LORY P., NARGEOT J., RASSENDREN F.A. & TIAHO F. Properties of L-type Ca channels expressed in *Xenopus laevis* oocytes by injection of rat heart mRNA. (C) 24*P*

LORY P., NARGEOT J., RICHARD S. & TIAHO F. Bay K 8644 slows inactivation of cardiac Ca channel current via a cyclic AMP-dependent mechanism in single rat ventricular myocytes. (C) 25*P*

LOVICK T.A. Effect of 5-HT in the ventrolateral medulla on the pressor response and analgesia evoked by stimulation of the dorsal periaqueductal grey matter in anaesthetized rats. (C) 84*P*

LUCIO F.J. *see* HALL A.C., HARVEY C.M., HENDRY B.M. & LUCIO F.J. Effects of hydrostatic pressure and membrane cholesterol depletion on L-lysine transport in human erythrocytes. (C) 89*P*

LUKOWIAK K. *see* HAYDON P.G., LUKOWIAK K., MAN-SON-HING H.J. & ZORAN M.J. Mechanisms of synaptic plasticity: modulation of secretory apparatus in cultured neuronal somata of *Helisoma trivolvis*. (C) 111*P*

LUKOWIAK K. *see* HAYDON P.G., LUKOWIAK K., MAN-SON-HING H.J. & ZORAN M.J. Mechanisms of synaptic plasticity: Use of a giant soma–soma synapse in cultured neuronal somata of *Helisoma trivolvis*. (PC) 147*P*

LUNN E.R. *see* BROWN M.C., LUNN E.R. & PERRY V.H. The route taken by axons growing down an intact distal nerve stump in mice. (PC) 148*P*

LUSH D. *see* DEARNLEY D.P., GOMEZ S., HALL M., LUSH D., POPE R. & WATSON M.S. A computerized system for analysing lung function data in man. (D) 4*P*

LUTTMANN M.L. *see* DANTZIG J.A., GOLDMAN Y.E., LUTTMANN M.L., TRENTHAM D.R. & WOODWARD S.K.A. Binding of caged ATP diastereoisomers to rigor cross-bridges in glycerol-extracted fibres of rabbit psoas muscle. (C) 61*P*

LUXON L.M., MERTON P.A., MORTON H.B. & SECKER B. A rotating chair for investigation of vestibulo-ocular function in man. (D) 7*P*

LYNN J. *see* CADY E.B., JONES D.A., LYNN J. & NEWHAM D.J. Changes in force and intracellular metabolites during fatigue of human skeletal muscle. 311

MACDONALD G.J. *see* DUGGAN K.A., HAWLEY C.M., MACDONALD G.J. & SHELLEY S. Sodium depletion decreases hepatic metabolism of vasoactive intestinal peptide in the rabbit 251

MACEWAN D.J. *see* JOHNSON M.S., MACEWAN D.J., MITCHELL R. & THOMSON F.J. Lithium enhances the inhibitory effect of protein kinase C on LHRH-induced $^{45}Ca^{2+}$ influx into rat anterior pituitary tissue *in vitro*. (PC) 185*P*

MACEWAN D.J. *see* JOHNSON M.S., MACEWAN D.J. & MITCHELL R. Depolarization-induced $^{45}Ca^{2+}$ influx into rat anterior pituitary cells *in vitro* can be modulated in two ways by protein kinase C. (PC) 186*P*

MCGUIGAN J.A.S. *see* BLATTER L.A., BURI Arlette & MCGUIGAN J.A.S. Free intracellular magnesium concentration in isolated ferret ventricular muscle and in frog skeletal muscle measured with ion-selective microelectrodes containing the new magnesium sensor ETH 5214. (PC) 154*P*

MCHALE N.G. & THORNBURY K.D. Increased leucocyte output from the popliteal node in response to stimulation of the sympathetic chain in the anaesthetized sheep. (PC) 176*P*

MACHEREY H.-J., PETERSEN K.-U., SPRAKTIES G. & WINTERHAGER J.-M. Effects of putative Cl channel blockers on electroneutral and electrogenic HCO_3 secretion by guinea-pig gallbladder epithelium *in vitro*. (C) 22*P*

MACKIE G.O. & MEECH R.W. Potassium channel family in the axons of jellyfish *Aglantha digitale*. (C) 14*P*

MCKINLEY Michael J. *see* CHODOBSKI Adam & MCKINLEY Michael J. Cerebral regulation of renal sodium excretion in sheep infused intravenously with hypertonic NaCl. 273

MACLENNAN D.H. *see* BIGELOW D.J., CLARKE D.M., INESI G., LOO T.W., MACLENNAN D.H. & SQUIER T.C. Correlation of function and molecular structure in the calcium transport ATPase of sarcoplasmic reticulum isolated from rabbit skeletal muscle. (C) 145*P*

MCMAHON B.R. *see* BOURNE G.B. & MCMAHON B.R. Control of cardiac output and its distribution in crustacean open circulatory systems. (C) 134*P*

MACONOCHIE D.J. & MARTIN J.L. The direct fitting of a differential equation to exponential data. (D) 9*P*

MADRID J.A. *see* CAMELLO P.J., FRANCIS L.P., MADRID J.A., PARIENTE J.A., SALIDO G.M. & SINGH J. Phorbol ester stimulates juice secretion and inhibits secretin-evoked secretory response in the pancreas of the anaesthetized rat. (C) 99*P*

MADRID J.A., POZO M.J. & SALIDO G.M. Cholecystokinin evokes contraction of gallbladder and sphincter of Oddi in the anaesthetized dog. (C) 100*P*

MAEDA Y. *see* ASHLEY C.C., BAGNI A., CECCHI G., GRIFFITHS P.J. & MAEDA Y. Time-resolved X-ray diffraction studies on intact single muscle fibres of *Rana temporaria*. (C) 58*P*

MAHAUT-SMITH M.P. Activation of Ca-dependent K channels in rat thymic lymphocytes by a mitogenic lectin. (C) 94*P*

MAN-SON-HING H.J. *see* HAYDON P.G., LUKOWIAK K., MAN-SON-HING H.J. & ZORAN M.J. Mechanisms of synaptic plasticity: modulation of secretory apparatus in cultured neuronal somata of *Helisoma trivolvis*. (C) 111*P*

MAN-SON-HING H.J. *see* HAYDON P.G., LUKOWIAK K., MAN-SON-HING H.J. & ZORAN M.J. Mechanisms of synaptic plasticity: Use of a giant soma–soma synapse in cultured neuronal somata of *Helisoma trivolvis*. (PC) 147*P*

MÅNSSON B. *see* EKSTRÖM J., GARRETT J.R., MÅNSSON B. & PROCTOR G.B. Changes in protein composition of parotid saliva from anaesthetized rats after chronic sympathectomy, assessed by hydrophobic interaction chromatography. (C) 97*P*

MARCUSSEN B. *see* KAWAI M. & MARCUSSEN B. Kinetics of MgATP binding, oscillatory power production, and the rate-limiting step in isolated insect muscle fibres. (C) 64*P*

MARKIEWICZ M. *see* BARKER P.M., MARKIEWICZ M., PARKER K.A., STRANG L.B. & WALTERS D.V. Induction of the adrenaline-dependent reabsorption of lung liquid in the fetal sheep by synergistic action of triiodothyronine and hydrocortisone. (C) 138*P*

MARTIN J.L. *see* MACONOCHIE D.J. & MARTIN J.L. The direct fitting of a differential equation to exponential data. (D) 9*P*

MARTIN K.A.C. *see* BERMAN N.J., DOUGLAS R.J. & MARTIN K.A.C. The conductances associated with inhibitory postsynaptic potentials are larger in visual cortical neurones *in vitro* than in similar neurones in intact, anaesthetized rats. (C) 107*P*

MATHIE A. *see* CULL-CANDY S.G., MATHIE A., SYMONDS C.J. & WYLLIE D.J.A. Distribution of quisqualate and kainate receptors in rat type-2 astrocytes and their progenitor cells in culture. (PC) 195*P*

MATSUDA H., MATSUURA H. & NOMA A. Erratum: Journal of Physiology 413 (1989) 441

MATSUURA H. *see* MATSUDA H., MATSUURA H. & NOMA A. Erratum: Journal of Physiology 413 (1989) 441

MATTHEWS B. *see* BOISSONADE F.M. & MATTHEWS B. The effects of anaesthesia on the properties of neurones in the trigeminal brain-stem nuclei of the cat. (C) 115*P*

MATTHEWS Gary, NEHER Erwin & PENNER Reinhold Second messenger-activated calcium influx in rat peritoneal mast cells 105

MATTHEWS Gary, NEHER Erwin & PENNER Reinhold Chloride conductance activated by external agonists and internal messengers in rat peritoneal mast cells. 131

MATTHEWS H.R. A laboratory-instrument controller based on a single-chip microcomputer. (D) 2*P*

MATTHEWS Margaret R. *see* CLOWRY G.J. & MATTHEWS Margaret R. The effects of innervation of an inappropriate target, and exogenous ganglioside treatment, upon the retrograde axonal transport of HRP in axotomized hypoglossal neurones in the rat. (C) 73*P*

MAZDA Genie Y., NISTRI A. & SIVILOTTI Lucia Inhibition of GABA-induced responses by Cl^- transport inhibitors in the frog optic tectum *in vitro*. (PC) 152*P*

MEECH R.W. *see* MACKIE G.O. & MEECH R.W. Potassium channel family in the axons of jellyfish *Aglantha digitale*. (C) 14*P*

MENINI A. *see* COLAMARTINO G., MENINI A. & TORRE V. The blocking effect of divalent cations on the cyclic GMP-activated current in excised patches from retinal rods of the tiger salamander. (C) 122*P*

MENINI Anna The effect of hydrogen ions on the cyclic GMP-activated currents carried by Na^+, K^+ or Li^+ in excised patches from retinal rods of the salamander. (C) 123*P*

MERTON P.A. *see* LUXON L.M., MERTON P.A., MORTON H.B. & SECKER B. A rotating chair for investigation of vestibulo-ocular function in man. (D) 7*P*

METZGER Joseph M. & MOSS Richard L. Effects on tension and stiffness due to reduced pH in isolated rabbit fast- and slow-twitch single skeletal muscle fibres. (C) 67*P*

MEVES H., POHL J.-A. & RUBLY N. A slow component in the gating current on-response of the frog node of Ranvier. (C) 15*P*

MEYER D.J. Jr *see* HUXLEY Virginia H. & MEYER D.J. Jr Volume status dependence of atrial peptide-induced changes in capillary water conductance: evidence from individually perfused mesenteric microvessels of decerebrate leopard frogs. (C) 78*P*

MILLAR N.C. *see* HOMSHER E. & MILLAR N.C. Tension transients induced by photolysis of caged phosphate in glycerinated rabbit soleus muscle fibres. (C) 62*P*

MITCHELL M.A. *see* BROWN V., CARLISLE A.J. & MITCHELL M.A. Avian plasma concentrations of zinc; responses to stressors in the domestic fowl (*Gallus domesticus*). (C) 129*P*

MITCHELL R. *see* JOHNSON M.S., MACEWAN D.J., MITCHELL R. & THOMSON F.J. Lithium enhances the inhibitory effect of protein kinase C on LHRH-induced $^{45}Ca^{2+}$ influx into rat anterior pituitary tissue *in vitro*. (PC) 185*P*

MITCHELL R. *see* JOHNSON M.S., MACEWAN D.J. & MITCHELL R. Depolarization-induced $^{45}Ca^{2+}$ influx into rat anterior pituitary cells *in vitro* can be modulated in two ways by protein kinase C. (PC) 186*P*

MOCHIDA S. *see* DAUZENROTH M-E., DOLLY J.O., HABERMANN E., MOCHIDA S., POULAIN B., TAUC L., WADSWORTH J.D.F. & WELLER U. Light chain of tetanus toxin can be internalized into a cholinergic synapse of *Aplysia* by the heavy chain of botulinum neurotoxin and depresses the transmitter release. (C) 71*P*

MOLL A. *see* CADY E.B., ELSHOVE H., JONES D.A. & MOLL A. The metabolic causes of slow relaxation in fatigued human skeletal muscle. 327

MONTGOMERY B.S.I. *see* BARO I., EISNER D.A., FRY C.H., LISTON T.G., MONTGOMERY B.S.I. & RAIMBACH S.J. Measurement of intracellular pH in isolated ferret detrusor smooth muscle cells. (C) 124*P*

MORAD M. *see* AGUS Z.S., DUKES I.D. & MORAD M. Divalent cations modulate transient outward current in isolated rat ventricular myocytes. (C) 28*P*

MORAD Martin *see* DUKES Iain D. & MORAD Martin Tedisamil increases the rate of inactivation of the transient outward current in single rat ventricular myocytes. (C) 33*P*

MORAN J. *see* BARKLEY G.L., GARDNER-MEDWIN A.R., MORAN J., NAGEL-LEIBY S., SIMKINS R.T., TEPLEY N. & WELCH K.M.A. Magnetic fields during spreading depression in the cortex of anaesthetized rabbits. (C) 101*P*

MOROVAT A. *see* DAUNCEY M.J. & MOROVAT A. Cold-induced changes in nuclear 3,5,3′-triiodothyronine receptors of porcine skeletal muscle. (PC) 174*P*

MORRISON J.F.B. *see* GUNASENA Karen T., MORRISON J.F.B., NIMMO A.J. & WHITAKER Elaine M. The effect of oestradiol and progesterone on neurokinin receptors in rat uterus. (C) 75*P*

MORTON H.B. *see* LUXON L.M., MERTON P.A., MORTON H.B. & SECKER B. A rotating chair for investigation of vestibulo-ocular function in man. (D) 7*P*

MOSHI M.J., PARRATT J.R., WAINWRIGHT C.L. & ZEITLIN I.J. Acid optimum kinin-forming enzymes in dog cardiovascular tissues. (PC) 177*P*

Moss Richard L. *see* METZGER Joseph M. & Moss Richard L. Effects on tension and stiffness due to reduced pH in isolated rabbit fast- and slow-twitch single skeletal muscle fibres. (C) 67*P*

MUÑOZ D.J.B. *see* BULLIVANT E.M.A. & MUÑOZ D.J.B. A natriuretic substance is released from the kidney in response to low rates of saline infusion into the renal artery of the anaesthetized rat. (C) 141*P*

MURPHY C. *see* BUCH J., COATS A., CONWAY J., MURPHY C. & SLEIGHT P. The baroreflex control of cardiac function in man. (C) 83*P*

MURRELL R.D. *see* BRAUN M., HAYDON D.A. & MURRELL R.D. The effects of *n*-alcohols on the nicotinic acetylcholine receptor channel in cultured rat myotubes. (C) 37*P*

NAGEL-LEIBY S. *see* BARKLEY G.L., GARDNER-MEDWIN A.R., MORAN J., NAGEL-LEIBY S., SIMKINS R.T., TEPLEY N. & WELCH K.M.A. Magnetic fields during spreading depression in the cortex of anaesthetized rabbits. (C) 101*P*

NAIRN Angus C. *see* BAHINSKI Anthony, GADSBY David C., GREENGARD Paul & NAIRN Angus C. Chloride conductance regulated by protein kinase A in isolated guinea-pig ventricular myocytes. (C) 32*P*

NARGEOT J. *see* LORY P., NARGEOT J., RASSENDREN F.A. & TIAHO F. Properties of L-type Ca channels expressed in *Xenopus laevis* oocytes by injection of rat heart mRNA. (C) 24*P*

NARGEOT J. *see* LORY P., NARGEOT J., RICHARD S. & TIAHO F. Bay K 8644 slows inactivation of cardiac Ca channel current via a cyclic AMP-dependent mechanism in single rat ventricular myocytes. (C) 25*P*

NEHER Erwin *see* MATTHEWS Gary, NEHER Erwin & PENNER Reinhold Second messenger-activated calcium influx in rat peritoneal mast cells 105

NEHER Erwin *see* MATTHEWS Gary, NEHER Erwin & PENNER Reinhold Chloride conductance activated by external agonists and internal messengers in rat peritoneal mast cells. 131

NEUBER-HESS M. & ROSE P.K. Morphology of axon terminals on proximal and distal dendrites of neck motoneurones in the anaesthetized cat. (C) 117*P*

NEUHAUS R. *see* CACHELIN A.B. & NEUHAUS R. The effect of Mg^{2+} on currents through single acetylcholine channels of rat PC12 cells. (C) 38*P*

NEWHAM D.J. *see* CADY E.B., JONES D.A., LYNN J. & NEWHAM D.J. Changes in force and intracellular metabolites during fatigue of human skeletal muscle. 311

NIMMO A.J. *see* GUNASENA Karen T., MORRISON J.F.B., NIMMO A.J. & WHITAKER Elaine M. The effect of oestradiol and progesterone on neurokinin receptors in rat uterus. (C) 75*P*

NISTRI A., SIVILOTTI Lucia & WELSH Diana M. Effects of NMDA and its antagonists on excitatory synaptic transmission in the frog optic tectum *in vitro*. (PC) 151*P*

NISTRI A. *see* MAZDA Genie Y., NISTRI A. & SIVILOTTI Lucia Inhibition of GABA-induced responses by Cl^- transport inhibitors in the frog optic tectum *in vitro*. (PC) 152*P*

NOIREAUD J. *see* LEOTY C. & NOIREAUD J. Two components of potassium contractures in isolated ferret cremaster fibres. (PC) 172*P*

NOMA A. *see* MATSUDA H., MATSUURA H. & NOMA A. Erratum: Journal of Physiology 413 (1989) 441

OJEDA C. *see* BERNENGO J.-C., LEMTIRI-CHLIEH F., OJEDA C. & PLATONOFF N. Measurement of cytosolic free Ca in cultured rat neonate and adult guinea-pig ventricular cardiac myocytes. (C) 45*P*

OJEDA C. *see* CHRISTÉ G., DE LA CHAPELLE B., LEMTIRI-CHLIEH F. & OJEDA C. Presence of a Cs-resistant transient outward current which distorts I_{Ca} measurements in guinea-pig single ventricular cells. (PC) 169*P*

OLDENBOURG Rudolf *see* INOUÉ Shinya & OLDENBOURG Rudolf Polarized light microscopy of striated muscle and analysis of edge birefringence. (C) 56*P*

O'NEILL S.C. *see* DONOSO P., EISNER D.A. & O'NEILL S.C. Measurement of intracellular $[Na^+]$ in isolated rat cardiac myocytes using the fluorescent indicator SBFI. (C) 48*P*

ORLOWSKI S. *see* BERTHON B., CHAMPEIL P., CLARET M., COMBETTES L., DOUCET E. & ORLOWSKI S. Fast kinetics of calcium release induced by myo-inositoltrisphosphate in permeabilized rat hepatocytes. (PC) 178*P*

ORTEGA Alicia *see* GONZALEZ-SERRATOS H., INESI G. & ORTEGA Alicia Effects of D-600 on isolated fibres and isolated sarcoplasmic reticulum of frog skeletal muscle. (C) 119*P*

OTUN H. *see* DUNLOP W., GILLESPIE J.I., GREENWELL J.R. & OTUN H. Measurements of intracellular calcium (Ca_i^{2+}) and sodium (Na_i^+) in single cultured rat smooth muscle cells (A10) using the fluorescent dyes Fura-2 and SBFI. (PC) 201*P*

PARIENTE J.A. *see* CAMELLO P.J., FRANCIS L.P., MADRID J.A., PARIENTE J.A., SALIDO G.M. & SINGH J. Phorbol ester stimulates juice secretion and inhibits secretin-evoked secretory response in the pancreas of the anaesthetized rat. (C) 99*P*

PARKER K.A. *see* BARKER P.M., MARKIEWICZ M., PARKER K.A., STRANG L.B. & WALTERS D.V. Induction of the adrenaline-dependent reabsorption of lung liquid in the fetal sheep by synergistic action of triiodothyronine and hydrocortisone. (C) 138*P*

PARRATT J.R. *see* MOSHI M.J., PARRATT J.R., WAINWRIGHT C.L. & ZEITLIN I.J. Acid optimum kinin-forming enzymes in dog cardiovascular tissues. (PC) 177*P*

PARRI H.R. *see* HOLDEN-DYE L., PARRI H.R. & WALKER R.J. Anion selectivity of the GABA-operated channel in somatic muscle cells of the parasitic nematode *Ascaris suum*. (PC) 194*P*

PATERSON Christopher A. *see* DELAMERE Nicholas A. & PATERSON Christopher A. Studies on calcium transport in a rabbit lens membrane vesicle preparation. (PC) 181*P*

PATON J.F.R., SILVA-CARVALHO L., SPYER K.M. & THOMPSON C.S. Role of the nucleus tractus solitarius in mediating the parabrachial pressor response in the decerebrate rabbit. (C) 86*P*

PAWSON Peter A. *see* GRINNELL Alan D. & PAWSON Peter A. Dependence of spontaneous release at frog junctions on synaptic strength, external calcium and terminal length. 397

PEACHEY L.D. *see* HUANG C.L.-H. & PEACHEY L.D. Charge movements during a frog skeletal muscle action potential. (C) 121*P*

PECKHAM M., SPARROW J.C. & WHITE D.C.S. Mechanical responses of skinned flight muscle from mutants of *Drosophila melanogaster* with abnormal amounts of actin or myosin. (PC) 160*P*

PENINGTON N.J. *see* KELLY J.S. & PENINGTON N.J. 5-Hydroxytryptamine inhibits the voltage-dependent calcium current of acutely dissociated central neurones from the adult rat dorsal raphe nucleus. (C) 35*P*

PENNER Reinhold *see* MATTHEWS Gary, NEHER Erwin & PENNER Reinhold Second messenger-activated calcium influx in rat peritoneal mast cells 105

PENNER Reinhold *see* MATTHEWS Gary, NEHER Erwin & PENNER Reinhold Chloride conductance activated by external agonists and internal messengers in rat peritoneal mast cells. 131

PERKINS M.N. *see* CAMPBELL E.A., DRAY A., PERKINS M.N. & SHAW W. Analgesic and anti-inflammatory activity of resiniferatoxin in the mouse and rat: comparison with capsaicin. (PC) 149*P*

PERRY V.H. *see* BROWN M.C., LUNN E.R. & PERRY V.H. The route taken by axons growing down an intact distal nerve stump in mice. (PC) 148*P*

PETERSEN K.-U. *see* MACHEREY H.-J., PETERSEN K.-U., SPRAKTIES G. & WINTERHAGER J.-M. Effects

of putative Cl channel blockers on electroneutral and electrogenic HCO_3 secretion by guinea-pig gallbladder epithelium *in vitro*. (C) 22*P*

PETERSEN O.H. *see* DUNNE M.J., GALLACHER D.V., PETERSEN O.H. & YULE D.I. The electrogenic sodium–alanine co-transport system in RINm5F insulin-secreting cells: patch-clamp and single cell $[Ca^{2+}]_i$ measurements. (C) 96*P*

PETRICCIONE DI VADI P. *see* GALLAGHER J., GARLICK P., HAMANN W., KEEVIL S., PETRICCIONE DI VADI P., PORTER D. & WEDLEY J.R. The use of magnetic resonance spectroscopy in assessing the efficacy of chemical lumbar sympathectomy in ischaemic vascular disease. (D) 3*P*

PHILLIS J.W. & WALTER G.A. Hypoxaemia-evoked release of inhibitory and excitatory amino acids from the superfused cerebral cortex of the anaesthetized rat. (C) 110*P*

PIAZZESI G. *see* LOMBARDI V. & PIAZZESI G. Mechanical properties of tetanized frog muscle fibres during the transitory phase of force response to lengthening. (C) 65*P*

PLANT T.D. Effects of 1,2-dioctanoyl-*sn*-glycerol (DiC_8) on ionic currents in cultured mouse pancreatic B-cells. (C) 21*P*

PLATONOFF N. *see* BERNENGO J.-C., LEMTIRI-CHLIEH F., OJEDA C. & PLATONOFF N. Measurement of cytosolic free Ca in cultured rat neonate and adult guinea-pig ventricular cardiac myocytes. (C) 45*P*

POGGESI G. *see* CECCHI G., COLOMO F., POGGESI G. & TESI C. A simple method to measure isometric twitch tension in single frog cardiac cells. (C) 49*P*

POGGIOLI Josiane & RENARD Dominique Effect of protein kinase C activation on GTP-mediated activation of a Ca-dependent phosphoinositidase C in rat heart. (C) 46*P*

POHL J.-A. *see* MEVES H., POHL J.-A. & RUBLY N. A slow component in the gating current on-response of the frog node of Ranvier. (C) 15*P*

POPE R. *see* DEARNLEY D.P., GOMEZ S., HALL M., LUSH D., POPE R. & WATSON M.S. A computerized system for analysing lung function data in man. (D) 4*P*

POPP David *see* GAMBLE Theresa S., POPP David & STEWART Alex A.E. Time-resolved X-ray diffraction measurements of the isolated ABRM from *Mytilus edulis*. (PC) 204*P*

PORTER D. *see* GALLAGHER J., GARLICK P., HAMANN W., KEEVIL S., PETRICCIONE DI VADI P., PORTER D. & WEDLEY J.R. The use of magnetic resonance spectroscopy in assessing the efficacy of chemical lumbar sympathectomy in ischaemic vascular disease. (D) 3*P*

POULAIN B. *see* DAUZENROTH M-E., DOLLY J.O., HABERMANN E., MOCHIDA S., POULAIN B., TAUC L., WADSWORTH J.D.F. & WELLER U. Light chain of tetanus toxin can be internalized into a cholinergic synapse of *Aplysia* by the heavy chain of botulinum neurotoxin and depresses the transmitter release. (C) 71*P*

POWIS D.A. Sodium–calcium exchange in cultured bovine chromaffin cells. (PC) 182*P*

POZO M.J. *see* MADRID J.A., POZO M.J. & SALIDO G.M. Cholecystokinin evokes contraction of gallbladder and sphincter of Oddi in the anaesthetized dog. (C) 100*P*

PRINS M. *see* KROESE A.B.A., PRINS M. & SCHELLART N.A.M. Regional differences in conduction velocity and fibre diameter in posterior lateral line afferent axons in the trout. (C) 136*P*

PRIST R. *see* ACATAUASSÚ-NUNES R., ALBUQUERQUE A.M.T., PRIST R. & ROCHA E SILVA M. Haemodynamics of cardiac tamponade after hypertonic NaCl (7·5%) Dextran-70 (6%) – HSD – injections in anaesthetized dogs. (C) 81*P*

PROCHAZKA A. *see* LLEWELLYN M., PROCHAZKA A. & YANG J.F. Human H-reflexes are reduced in difficult beam-walking compared to stance and normal walking. (C) 104*P*

PROCHAZKA A. *see* LLEWELLYN M., PROCHAZKA A. & TREND P. Control of ankle extensor activation prior to and after foot contact in cat step cycle. (PC) 146*P*

PROCTOR G.B. *see* EKSTRÖM J., GARRETT J.R., MÅNSSON B. & PROCTOR G.B. Changes in protein composition of parotid saliva from anaesthetized rats after chronic sympathectomy, assessed by hydrophobic interaction chromatography. (C) 97*P*

RAIMBACH S.J. *see* BARO I., EISNER D.A., FRY C.H., LISTON T.G., MONTGOMERY B.S.I. & RAIMBACH S.J. Measurement of intracellular pH in isolated ferret detrusor smooth muscle cells. (C) 124*P*

RALL J.A. *see* HOU T.-t., JOHNSON J.D. & RALL J.A. Mg^{2+} and Ca^{2+} dissociation rates from frog parvalbumin at 0 °C: implications for role of parvalbumin in skeletal muscle relaxation. (C) 120*P*

RANATUNGA K.W. & THOMAS P.E. Differences in the extent of transformation in cross-reinnervated rat soleus muscle. (C) 53*P*

RANATUNGA K.W. *see* FORTUNE N.S., GEEVES M.A. & RANATUNGA K.W. Tension transients initiated by pressure perturbation in isolated rabbit skinned muscle fibres. (PC) 158*P*

RANDALL D.J. *see* AOTA S. & RANDALL D.J. Catecholamines and the central control of breathing in fish. (C) 130*P*

RASSENDREN F.A. *see* LORY P., NARGEOT J., RASSENDREN F.A. & TIAHO F. Properties of L-type Ca channels expressed in *Xenopus laevis* oocytes by injection of rat heart mRNA. (C) 24*P*

REAST Rebecca & SAGE Stewart O. Evidence for biphasic calcium influx evoked by ADP in Fura-2-loaded human platelets. (C) 93*P*

REEH P.W. *see* KIRCHHOFF C.G., REEH P.W. & WADDELL P.J. Sensory endings of C- and A-fibres are differentially sensitive to tetrodotoxin in the rat skin *in vitro*. (C) 116*P*

RENARD Dominique *see* POGGIOLI Josiane & RENARD Dominique Effect of protein kinase C activation on GTP-mediated activation of a Ca-dependent phosphoinositidase C in rat heart. (C) 46*P*

RICHARD S. *see* LORY P., NARGEOT J., RICHARD S. & TIAHO F. Bay K 8644 slows inactivation of cardiac Ca channel current via a cyclic AMP-dependent mechanism in single rat ventricular myocytes. (C) 25*P*

RIDGE R.M.A.P. *see* BETZ W.J., CHUA M. & RIDGE R.M.A.P. Inhibitory interactions between motoneurone terminals in neonatal rat lumbrical muscle. 25

ROBERTS C. *see* ANDERSON L.S., BUTLER P.J., ROBERTS C., SMALE K., SNOW D.H. & WOAKES A.J. Partial pressures of arterial and end-tidal gases during graded exercise in thoroughbred race-horses. (C) 131*P*

ROCHA E SILVA M. *see* ACATAUASSÚ-NUNES R., ALBUQUERQUE A.M.T., PRIST R. & ROCHA E SILVA M. Haemodynamics of cardiac tamponade after hypertonic NaCl (7·5%) Dextran-70 (6%) – HSD – injections in anaesthetized dogs. (C) 81*P*

ROCHE S.W. *see* FARMER S.F., INGRAM D.A., ROCHE S.W. & STEPHENS J.A. Evidence that mirror movements in the Klippel–Feil syndrome in man are produced by activity in branched-stem presynaptic inputs from corticospinal neurones. (C) 102*P*

ROOS K.P. & TAYLOR S.R. Striation dynamics in isolated rat cardiac myocytes revealed by computer vision. (C) 50*P*

ROSE P.K. *see* NEUBER-HESS M. & ROSE P.K. Morphology of axon terminals on proximal and distal dendrites of neck motoneurones in the anaesthetized cat. (C) 117*P*

ROSENBLATT J. David Return and subsequent diminution of the relative contractile force in rat extensor digitorum longus muscles following bupivacaine injection. (PC) 159*P*

ROWLERSON Anthea M. *see* BLACKSHAW Susanne E. & ROWLERSON Anthea M. ATPase activity and antibodies to vertebrate slow myosins distinguish fibre types in leech body wall muscle. (C) 51*P*

RUBINSTEIN Chaim T. & SHRAGER Peter Optical recording of action potentials in demyelinated and remyelinating frog nerve fibres. (C) 18*P*

RUBLY N. *see* MEVES H., POHL J.-A. & RUBLY N. A slow component in the gating current on-response of the frog node of Ranvier. (C) 15*P*

RUDD B.T. *see* DAUNCEY M.J., HOLDER G., INGRAM D.L., RUDD B.T. & SHAKESPEAR R.A. Thermal and nutritional regulation of insulin-like growth factor-I and cortisol in the young pig. (PC) 175*P*

RYDER Kathryn O. *see* BRYANT S.M., HART G. & RYDER Kathryn O. Captopril reduces L-type calcium current in isolated guinea-pig ventricular myocytes. (PC) 167*P*

SAGE S.O. *see* ADAMS D.J., VAN BREEMEN C., CANNELL M.B. & SAGE S.O. Na:Ca exchange in Fura-2-loaded bovine pulmonary artery endothelial cell monolayers. (PC) 183*P*

SAGE Stewart O. *see* REAST Rebecca & SAGE Stewart O. Evidence for biphasic calcium influx evoked by ADP in Fura-2-loaded human platelets. (C) 93*P*

SALIDO G.M. *see* CAMELLO P.J., FRANCIS L.P., MADRID J.A., PARIENTE J.A., SALIDO G.M. & SINGH J. Phorbol ester stimulates juice secretion and inhibits secretin-evoked secretory response in the pancreas of the anaesthetized rat. (C) 99*P*

SALIDO G.M. *see* MADRID J.A., POZO M.J. & SALIDO G.M. Cholecystokinin evokes contraction of gallbladder and sphincter of Oddi in the anaesthetized dog. (C) 100*P*

SALT T.E. & SILLITO A.M. The NMDA-receptor mediated component of the visual response of cells in the feline dorsal lateral geniculate nucleus (dLGN) in the absence of corticofugal feedback. (C) 108*P*

SALT T.E. Effects of cortical inactivation on responses of ventrobasal thalamus (VB) neurones to somatosensory stimuli in the anaesthetized rat. (C) 113*P*

SANDERS K.M. *see* BURKE E.P. & SANDERS K.M. Sodium-pump current in canine colonic myocytes is sufficient to provide a large contribution to membrane potential. (PC) 171*P*

SANSOM M.S.P. *see* BALL F.G., SANSOM M.S.P. & USHERWOOD P.N.R. A comparison of different gating models for voltage- and receptor-gated ion channels. (PC) 189*P*

SATTELLE D.B. *see* BUCKINGHAM S.D., HARRISON J.B., LEECH C.A. & SATTELLE D.B. Nicotinic acetylcholine receptor activation in dissociated neurones from *Periplaneta americana* and *Musca domestica*. (PC) 193*P*

SAUNDERS N.R. *see* AIKEN S.P., HORN N.M. & SAUNDERS N.R. Effect of histidine on zinc distribution in anaesthetized rats. (C) 144*P*

SCHELLART N.A.M. *see* KROESE A.B.A., PRINS M. & SCHELLART N.A.M. Regional differences in conduction velocity and fibre diameter in posterior lateral line afferent axons in the trout. (C) 136*P*

SCHLUE W.-R. *see* BORON W.F., BOYARSKY G., GANZ M.B. & SCHLUE W.-R. The effect of DIDS on pH regulation of isolated single rat mesangial cells examined in the absence of HCO_3^-. (C) 125*P*

SCHWIENING Christof J. & THOMAS Roger C. Mechanism of intracellular pH (pH_i) regulation by isolated locust neurones. (C) 126*P*

SCOTT R.H. *see* ADAMS M.E., BINDOKAS V.P., DOLPHIN A.C. & SCOTT R.H. The inhibition of Ca^{2+} channel currents in cultured neonatal rat dorsal root ganglion (DRG) neurones by ω-Agatoxin-1A (a funnel web spider toxin). (C) 34*P*

SEARS T.A. *see* KIRKWOOD P.A. & SEARS T.A. Field potentials in the contralateral ventral horn evoked by individual thoracic respiratory interneurones in the anaesthetized cat. (C) 106*P*

SECKER B. *see* LUXON L.M., MERTON P.A., MORTON H.B. & SECKER B. A rotating chair for investigation of vestibulo-ocular function in man. (D) 7*P*

SEO Y. *see* CASE R.M., SEO Y. & STEWARD M.C. Regulation of intracellular pH in the isolated, perfused rabbit mandibular salivary gland studied by ^{31}P NMR spectroscopy. (C) 127*P*

SEPÚLVEDA F.V. *see* GIRALDEZ F., SEPÚLVEDA F.V., SHEPPARD D.N. & VALVERDE M.A. Potassium currents in isolated *Necturus* enterocytes. (C) 13*P*

SHAFI Shahida *see* BORN G.V.R. & SHAFI Shahida Evidence against the atherogenic passage of low-density lipoproteins between vascular endothelial cells in conscious rabbits. (C) 80*P*

SHAKESPEAR R.A. *see* DAUNCEY M.J., HOLDER G., INGRAM D.L., RUDD B.T. & SHAKESPEAR R.A. Thermal and nutritional regulation of insulin-like growth factor-I and cortisol in the young pig. (PC) 175*P*

SHAW W. *see* CAMPBELL E.A., DRAY A., PERKINS M.N. & SHAW W. Analgesic and anti-inflammatory activity of resiniferatoxin in the mouse and rat: comparison with capsaicin. (PC) 149*P*

SHELLEY S. *see* DUGGAN K.A., HAWLEY C.M., MACDONALD G.J. & SHELLEY S. Sodium depletion decreases hepatic metabolism of vasoactive intestinal peptide in the rabbit 251

SHEN S. *see* BASKIN R.J., SHEN S. & YEH Y. Optical measurements on isolated single fibres from insect flight muscle. (C) 55*P*

SHEPPARD D.N. *see* GIRALDEZ F., SEPÚLVEDA F.V., SHEPPARD D.N. & VALVERDE M.A. Potassium currents in isolated *Necturus* enterocytes. (C) 13*P*

SHERIDAN D.J. *see* FLORES N.A., JUHASZ C., SHERIDAN D.J. & WINTERTON S.J. An IBM PC AT based system for the analysis of cardiac action potentials in real time. (D) 8*P*

SHIRLEY D.G., SKINNER J. & WALTER S.J. Effect of sodium depletion on renal lithium reabsorption in conscious rats. (C) 92*P*

SHRAGER Peter *see* RUBINSTEIN Chaim T. & SHRAGER Peter Optical recording of action potentials in demyelinated and remyelinating frog nerve fibres. (C) 18*P*

SHRIMPTON C. *see* IRVING M., KENDRICK-JONES J., SHRIMPTON C. & SLEEP J. The orientation of rhodamine probes attached to myosin light chain-2 (LC-2) in relaxing and rigor conditions in single fibres isolated from rabbit psoas muscle. (C) 57*P*

SILLITO A.M. *see* SALT T.E. & SILLITO A.M. The NMDA-receptor mediated component of the visual response of cells in the feline dorsal lateral geniculate nucleus (dLGN) in the absence of corticofugal feedback. (C) 108*P*

SILVA-CARVALHO L. *see* PATON J.F.R., SILVA-CARVALHO L., SPYER K.M. & THOMPSON C.S. Role of the nucleus tractus solitarius in mediating the parabrachial pressor response in the decerebrate rabbit. (C) 86*P*

SIMKINS R.T. *see* BARKLEY G.L., GARDNER-MEDWIN A.R., MORAN J., NAGEL-LEIBY S., SIMKINS R.T., TEPLEY N. & WELCH K.M.A. Magnetic fields during spreading depression in the cortex of anaesthetized rabbits. (C) 101*P*

SIMMONS R.M. *see* BRENNER B., FERENCZI M.A., IRVING M., KAPLAN J., SIMMONS R.M. & TOWNS-ANDREWS E. Structural changes upon activation of single isolated muscle fibres of the frog by photolysis of 'caged calcium' studied by time-resolved X-ray diffraction. (C) 59*P*

SINGH J. *see* CAMELLO P.J., FRANCIS L.P., MADRID J.A., PARIENTE J.A., SALIDO G.M. & SINGH J.

Phorbol ester stimulates juice secretion and inhibits secretin-evoked secretory response in the pancreas of the anaesthetized rat. (C) 99*P*

SIPIDO K.R. & WIER W.G. $[Ca^{2+}]_i$ transients in isolated guinea-pig ventricular myocytes in the absence of sodium–calcium (Na–Ca) exchange and sarcoplasmic reticulum function. (PC) 170*P*

SITSAPESAN R. & WILLIAMS A.J. Caffeine activation of single calcium-release channels from the sheep cardiac sarcoplasmic reticulum. (C) 26*P*

SIVILOTTI Lucia *see* NISTRI A., SIVILOTTI Lucia & WELSH Diana M. Effects of NMDA and its antagonists on excitatory synaptic transmission in the frog optic tectum *in vitro*. (PC) 151*P*

SIVILOTTI Lucia *see* MAZDA Genie Y., NISTRI A. & SIVILOTTI Lucia Inhibition of GABA-induced responses by Cl^- transport inhibitors in the frog optic tectum *in vitro*. (PC) 152*P*

SJÖGREN C. *see* ANDERSSON P.-O., SJÖGREN C., UVNÄS B. & UVNÄS-MOBERG K. The role of vasoactive intestinal polypeptide (VIP) in mediating responses of the urinary bladder to stimulation of the pelvic and hypogastric nerves in anaesthetized dogs. (C) 74*P*

SKINNER J. *see* SHIRLEY D.G., SKINNER J. & WALTER S.J. Effect of sodium depletion on renal lithium reabsorption in conscious rats. (C) 92*P*

SLEEP J. *see* IRVING M., KENDRICK-JONES J., SHRIMPTON C. & SLEEP J. The orientation of rhodamine probes attached to myosin light chain-2 (LC-2) in relaxing and rigor conditions in single fibres isolated from rabbit psoas muscle. (C) 57*P*

SLEIGHT P. *see* BUCH J., COATS A., CONWAY J., MURPHY C. & SLEIGHT P. The baroreflex control of cardiac function in man. (C) 83*P*

SLEVIN K. *see* JACKSON A.A., SLEVIN K. & WATERLOW J.C. A model of whole body protein turnover in man that includes a pool turning over by life-time kinetics. (C) 143*P*

SMALE K. *see* ANDERSON L.S., BUTLER P.J., ROBERTS C., SMALE K., SNOW D.H. & WOAKES A.J. Partial pressures of arterial and end-tidal gases during graded exercise in thoroughbred race-horses. (C) 131*P*

SMART T.G. *see* BOWIE D. & SMART T.G. Properties of a kainate receptor expressed in *Xenopus laevis* oocytes after injection with bovine brain mRNA. (PC) 196*P*

SMITH Allan *see* FERRELL William R. & SMITH Allan The effect of loading on position sense at the proximal interphalangeal joint of the human finger. 145

SMITH N.F. *see* EDDY F.B., GRIERSON C., HAZON N. & SMITH N.F. Effects of atrial natriuretic peptide (ANP) on circulatory and ion regulatory physiology of rainbow trout (*Salmo gairdneri* Richardson) fed normal or high levels of dietary salt. (C) 137*P*

SMOLLEY John R. *see* BRIDGE John H.B., SMOLLEY John R. & SPITZER Kenneth W. A discrete inward Na^+–Ca^{2+} exchange current accompanies the mechanical relaxation of isolated guinea-pig ventricular cells. (PC) 168*P*

SNOW D.H. *see* ANDERSON L.S., BUTLER P.J., ROBERTS C., SMALE K., SNOW D.H. & WOAKES A.J. Partial pressures of arterial and end-tidal gases during graded exercise in thoroughbred race-horses. (C) 131*P*

SO I.S. *see* EARM Y.E., HO W.K. & SO I.S. Time course of Na–Ca exchange current in isolated rabbit atrial cells. (C) 41*P*

SOLOMON A.K. *see* HARRIS H.W. Jr, JANOSHAZI A., KIKERI D., SOLOMON A.K. & ZEIDEL M.L. Proton permeability of toad bladder aggrephore membranes: role of the water channel in proton transport. (C) 145*P*

SOUTHAN A.P. & WANN K.T. High helium pressure modifies the repetitive discharge of CA1 pyramidal neurones in the rat hippocampus *in vitro*. (C) 17*P*

SPARROW J.C. *see* PECKHAM M., SPARROW J.C. & WHITE D.C.S. Mechanical responses of skinned flight muscle from mutants of *Drosophila melanogaster* with abnormal amounts of actin or myosin. (PC) 160*P*

SPITZER Kenneth W. *see* BRIDGE John H.B., SMOLLEY John R. & SPITZER Kenneth W. A discrete inward Na^+–Ca^{2+} exchange current accompanies the mechanical relaxation of isolated guinea-pig ventricular cells. (PC) 168*P*

SPRAKTIES G. *see* MACHEREY H.-J., PETERSEN K.-U., SPRAKTIES G. & WINTERHAGER J.-M. Effects of putative Cl channel blockers on electroneutral and electrogenic HCO_3 secretion by guinea-pig gallbladder epithelium *in vitro*. (C) 22*P*

SPRUCE A.E. *see* ALMERS W., IWATA A., SPRUCE A.E. & WHITE J.M. Patch clamp studies of fusion of human erythrocytes to murine fibroblasts mediated by influenza virus fusion protein. (C) 95*P*

SPYER K.M. *see* PATON J.F.R., SILVA-CARVALHO L., SPYER K.M. & THOMPSON C.S. Role of the nucleus tractus solitarius in mediating the parabrachial pressor response in the decerebrate rabbit. (C) 86*P*

SQUIER T.C. *see* BIGELOW D.J., CLARKE D.M., INESI G., LOO T.W., MACLENNAN D.H. & SQUIER T.C. Correlation of function and molecular structure in the calcium transport ATPase of sarcoplasmic reticulum isolated from rabbit skeletal muscle. (C) 145*P*

ST JOHN WALTER M. & ZHOU Dan Differing control of neural activities during various portions of expiration in the cat 189

STANDEN N.B. *see* DAVIES N.W., STANDEN N.B. & STANFIELD P.R. The kinetic behaviour of K_{ATP} channels in isolated membrane patches of frog skeletal muscle is regulated by internal pH. (PC) 188*P*

STANFIELD P.R. *see* DAVIES N.W., STANDEN N.B. & STANFIELD P.R. The kinetic behaviour of K_{ATP} channels in isolated membrane patches of frog skeletal muscle is regulated by internal pH. (PC) 188*P*

STEPHENS J.A. *see* EVANS A.L., HARRISON Linda M. & STEPHENS J.A. Task-dependent changes in cutaneous reflexes recorded from various muscles controlling finger movement in man. 1

STEPHENS J.A. *see* DATTA A.K., HARRISON Linda M. & STEPHENS J.A. Task-dependent changes in the size of response to magnetic brain stimulation in human first dorsal interosseous muscle. 13

STEPHENS J.A. *see* HARRISON Linda M., IRONTON R. & STEPHENS J.A. Cross correlation analysis of multi-unit EMG recordings in man. (D) 6*P*

STEPHENS J.A. *see* FARMER S.F., INGRAM D.A., ROCHE S.W. & STEPHENS J.A. Evidence that mirror movements in the Klippel–Feil syndrome in man are produced by activity in branched-stem presynaptic inputs from corticospinal neurones. (C) 102*P*

STEPHENSON D.G. *see* LAMB G.D. & STEPHENSON D.G. Activation and reduction of calcium release in skinned muscle fibres from the cane toad. (C) 70*P*

STEPHENSON D.G. & WEST J.M. Can the contractile apparatus in single muscle fibres from the extensor claw muscle of the yabby be fully activated by Sr^{2+} and Ca^{2+}? (PC) 162*P*

STEWARD M.C. *see* CASE R.M., SEO Y. & STEWARD M.C. Regulation of intracellular pH in the isolated, perfused rabbit mandibular salivary gland studied by ^{31}P NMR spectroscopy. (C) 127*P*

STEWART Alex A.E. *see* GAMBLE Theresa S., POPP David & STEWART Alex A.E. Time-resolved X-ray diffraction measurements of the isolated ABRM from *Mytilus edulis*. (PC) 204*P*

STRANG L.B. *see* BARKER P.M., MARKIEWICZ M., PARKER K.A., STRANG L.B. & WALTERS D.V. Induction of the adrenaline-dependent reabsorption of lung liquid in the fetal sheep by synergistic action of triiodothyronine and hydrocortisone. (C) 138*P*

SULEIMAN A.H. *see* ANDERSON L.C. & SULEIMAN A.H. Effects of streptozotocin-induced diabetes on sympathetically evoked submandibular salivary secretion in anaesthetized rats. (C) 98*P*

SULEIMAN M.S. *see* CHAPMAN R.A. & SULEIMAN M.S. The effect of hypothermia on $[Na]_i$ and tension development during the Ca paradox in isolated ferret ventricular muscle. (C) 47*P*

SULLIVAN S.K. *see* FIELD M., GRENSPAN N.R., SULLIVAN S.K. & SWAMY K. Phorbol myristate acetate (PMA) inactivates an epithelial potassium channel from flounder intestine expressed by *Xenopus* oocytes. (C) 23*P*

SWAMY K. *see* FIELD M., GRENSPAN N.R., SULLIVAN S.K. & SWAMY K. Phorbol myristate acetate (PMA) inactivates an epithelial potassium channel from flounder intestine expressed by *Xenopus* oocytes. (C) 23*P*

SYED N.I. & WINLOW W. An identified, high-order neurone driving the whole animal withdrawal response in *Lymnaea*. (C) 72*P*

SYMONDS C.J. *see* CULL-CANDY S.G., MATHIE A., SYMONDS C.J. & WYLLIE D.J.A. Distribution of quisqualate and kainate receptors in rat type-2 astrocytes and their progenitor cells in culture. (PC) 195*P*

TANSEY E.M. The scientific achievements of Lord Adrian, FRS (1899–1977). (D) 5*P*

TATTERSALL J.E.H. Acetylcholinesterase reactivators block open nicotinic ion channels in isolated adult mouse muscle fibres. (PC) 187*P*

TAUC L. *see* DAUZENROTH M-E., DOLLY J.O., HABERMANN E., MOCHIDA S., POULAIN B., TAUC L., WADSWORTH J.D.F. & WELLER U. Light chain of tetanus toxin can be internalized into a cholinergic synapse of *Aplysia* by the heavy chain of botulinum neurotoxin and depresses the transmitter release. (C) 71*P*

TAYLOR A. *see* GOMEZ K. & TAYLOR A. Spectral analysis of the demodulated EMG in isometric and isotonic human muscle contractions. (C) 105*P*

TAYLOR E.W. & WILSON R.W. The cardiovascular and respiratory responses to intra-arterial injection of adrenaline in the dogfish *Scyliorhinus canicula* L. (C) 133*P*

TAYLOR E.W. *see* GEE H. & TAYLOR E.W. Detection of sinus arrhythmia in the human fetus *in utero* using ultrasound and Doppler blood-flow waveforms in the major vessels. (PC) 184*P*

TAYLOR S.R. *see* ROOS K.P. & TAYLOR S.R. Striation dynamics in isolated rat cardiac myocytes revealed by computer vision. (C) 50*P*

TEPLEY N. *see* BARKLEY G.L., GARDNER-MEDWIN A.R., MORAN J., NAGEL-LEIBY S., SIMKINS R.T., TEPLEY N. & WELCH K.M.A. Magnetic fields during spreading depression in the cortex of anaesthetized rabbits. (C) 101*P*

TESI C. *see* CECCHI G., COLOMO F., POGGESI G. & TESI C. A simple method to measure isometric twitch tension in single frog cardiac cells. (C) 49*P*

THOMAS C.K. *see* BIGLAND-RITCHIE B., JOHANSSON R.S., THOMAS C.K. & WESTLING G. Twitch fusion in human motor units studied by motor axon stimulation. (C) 54*P*

THOMAS P.E. *see* RANATUNGA K.W. & THOMAS P.E. Differences in the extent of transformation in cross-reinnervated rat soleus muscle. (C) 53*P*

THOMAS Roger C. *see* SCHWIENING Christof J. & THOMAS Roger C. Mechanism of intracellular pH (pH_i) regulation by isolated locust neurones. (C) 126*P*

THOMPSON C.S. *see* PATON J.F.R., SILVA-CARVALHO L., SPYER K.M. & THOMPSON C.S. Role of the nucleus tractus solitarius in mediating the parabrachial pressor response in the decerebrate rabbit. (C) 86*P*

THOMSON F.J. *see* JOHNSON M.S., MACEWAN D.J., MITCHELL R. & THOMSON F.J. Lithium enhances the inhibitory effect of protein kinase C on LHRH-induced $^{45}Ca^{2+}$ influx into rat anterior pituitary tissue *in vitro*. (PC) 185*P*

THORNBURY K.D. *see* MCHALE N.G. & THORNBURY K.D. Increased leucocyte output from the popliteal node in response to stimulation of the sympathetic chain in the anaesthetized sheep. (PC) 176*P*

TIAHO F. *see* LORY P., NARGEOT J., RASSENDREN F.A. & TIAHO F. Properties of L-type Ca channels expressed in *Xenopus laevis* oocytes by injection of rat heart mRNA. (C) 24*P*

TIAHO F. *see* LORY P., NARGEOT J., RICHARD S. & TIAHO F. Bay K 8644 slows inactivation of cardiac Ca channel current via a cyclic AMP-dependent mechanism in single rat ventricular myocytes. (C) 25*P*

TOMLINSON K.C. *see* BENNETT T., GARDINER S.M. & TOMLINSON K.C. Cardiovascular and drinking responses to intracerebroventricular angiotensin II in streptozotocin-treated rats. (C) 82*P*

TORRE V. *see* COLAMARTINO G., MENINI A. & TORRE V. The blocking effect of divalent cations on the cyclic GMP-activated current in excised patches from retinal rods of the tiger salamander. (C) 122*P*

TOWNS-ANDREWS E. *see* BRENNER B., FERENCZI M.A., IRVING M., KAPLAN J., SIMMONS R.M. & TOWNS-ANDREWS E. Structural changes upon activation of single isolated muscle fibres of the frog by photolysis of 'caged calcium' studied by time-resolved X-ray diffraction. (C) 59*P*

TREND P. *see* LLEWELLYN M., PROCHAZKA A. & TREND P. Control of ankle extensor activation prior to and after foot contact in cat step cycle. (PC) 146*P*

TRENTHAM D.R. *see* DANTZIG J.A., GOLDMAN Y.E., LUTTMANN M.L., TRENTHAM D.R. & WOODWARD S.K.A. Binding of caged ATP diastereoisomers to rigor cross-bridges in glycerol-extracted fibres of rabbit psoas muscle. (C) 61*P*

TRENTHAM D.R. *see* FERENCZI M.A., GOLDMAN Y.E. & TRENTHAM D.R. Relaxation of permeabilized, isolated muscle fibres of the rabbit by rapid chelation of Ca^{2+}-ions through laser pulse photolysis of 'caged-BAPTA'. (PC) 155*P*

TROMBA C. *see* COHEN I. & TROMBA C. Inactivation of i_{K1} in isolated canine Purkinje myocytes is induced by isoprenaline but not blocked by propranolol. (C) 40*P*

TSAI T.D. *see* LEE K.S. & TSAI T.D. Open channel block of the fast inward sodium current by TTX in single monkey heart cells? (PC) 190*P*

TURNER M.R. Permeability of monolayers of cultured bovine endothelium in the presence and absence of bovine serum albumin. (C) 79*P*

USHERWOOD P.N.R. *see* BALL F.G., SANSOM M.S.P. & USHERWOOD P.N.R. A comparison of different gating models for voltage- and receptor-gated ion channels. (PC) 189*P*

UVNÄS B. *see* ANDERSSON P.-O., SJÖGREN C., UVNÄS B. & UVNÄS-MOBERG K. The role of vasoactive intestinal polypeptide (VIP) in mediating responses of the urinary bladder to stimulation of the pelvic and hypogastric nerves in anaesthetized dogs. (C) 74*P*

UVNÄS-MOBERG K. *see* ANDERSSON P.-O., SJÖGREN C., UVNÄS B. & UVNÄS-MOBERG K. The role of vasoactive intestinal polypeptide (VIP) in mediating responses of the urinary bladder to stimulation of the pelvic and hypogastric nerves in anaesthetized dogs. (C) 74*P*

VALVERDE M.A. *see* GIRALDEZ F., SEPÚLVEDA F.V., SHEPPARD D.N. & VALVERDE M.A. Potassium currents in isolated *Necturus* enterocytes. (C) 13*P*

VAN BREEMEN C. *see* ADAMS D.J., VAN BREEMEN C., CANNELL M.B. & SAGE S.O. Na:Ca exchange in Fura-2-loaded bovine pulmonary artery endothelial cell monolayers. (PC) 183*P*

VAN HELDEN D.F. Noradrenaline-induced transient depolarizations in the smooth muscle of isolated guinea-pig mesenteric lymphatics. (PC) 173*P*

VAN VLIET B.N. *see* BACKHOUSE S.S., HARPER A.A., VAN VLIET B.N. & WEST N.H. Localization and conduction velocities of arterial baroreceptors in the anaesthetized snake. (C) 135*P*

VAUGHAN-JONES R.D. *see* BOUNTRA C. & VAUGHAN-JONES R.D. Effect of intracellular and extracellular pH on contraction in isolated, mammalian cardiac muscle. 163

VAUGHAN-JONES R.D. *see* GASSER R. & VAUGHAN-JONES R.D. Myocardial ischaemia: release of potassium can be inhibited by tolbutamide and glibenclamide in isolated guinea-pig papillary muscle. (C) 43*P*

VAUGHAN-JONES R.D. & WU Mei-Lin Intracellular intrinsic buffering power varies with pH_i in the isolated sheep cardiac Purkinje fibre. (PC) 165*P*

WADDELL P.J. *see* KIRCHHOFF C.G., REEH P.W. & WADDELL P.J. Sensory endings of C- and A-fibres are differentially sensitive to tetrodotoxin in the rat skin *in vitro*. (C) 116*P*

WADSWORTH J.D.F. *see* DAUZENROTH M-E., DOLLY J.O., HABERMANN E., MOCHIDA S., POULAIN B., TAUC L., WADSWORTH J.D.F. & WELLER U. Light chain of tetanus toxin can be internalized into a cholinergic synapse of *Aplysia* by the heavy chain of botulinum neurotoxin and depresses the transmitter release. (C) 71*P*

WAINWRIGHT C.L. *see* MOSHI M.J., PARRATT J.R., WAINWRIGHT C.L. & ZEITLIN I.J. Acid optimum kinin-forming enzymes in dog cardiovascular tissues. (PC) 177*P*

WAKEFIELD Bruce T. *see* GAGE Peter W., LAMB Graham D. & WAKEFIELD Bruce T. Transient and persistent sodium currents in normal and denervated mammalian skeletal muscle. 427

WAKUI M. *see* KAO C.Y., WAKUI M., WANG S.Y. & YOSHINO M. The outward current of the isolated rat myometrium. (C) 20*P*

WALKER R.J. *see* HOLDEN-DYE L., PARRI H.R. & WALKER R.J. Anion selectivity of the GABA-operated channel in somatic muscle cells of the parasitic nematode *Ascaris suum*. (PC) 194*P*

WALTER G.A. *see* PHILLIS J.W. & WALTER G.A. Hypoxaemia-evoked release of inhibitory and excitatory amino acids from the superfused cerebral cortex of the anaesthetized rat. (C) 110*P*

WALTER S.J. *see* SHIRLEY D.G., SKINNER J. & WALTER S.J. Effect of sodium depletion on renal lithium reabsorption in conscious rats. (C) 92*P*

WALTERS D.V. *see* BARKER P.M., MARKIEWICZ M., PARKER K.A., STRANG L.B. & WALTERS D.V. Induction of the adrenaline-dependent reabsorption of lung liquid in the fetal sheep by synergistic action of triiodothyronine and hydrocortisone. (C) 138*P*

WALTON T. *see* CAVIERES J.D. & WALTON T. Pre-steady-state kinetics of product release by Na,K-ATPase purified from pig kidney. (C) 90*P*

WANG S.Y. *see* KAO C.Y., WAKUI M., WANG S.Y. & YOSHINO M. The outward current of the isolated rat myometrium. (C) 20*P*

WANN K.T. *see* SOUTHAN A.P. & WANN K.T. High helium pressure modifies the repetitive discharge of CA1 pyramidal neurones in the rat hippocampus *in vitro*. (C) 17*P*

WARD Christine E. *see* CLAFLIN Dennis R., FAULKNER John A. & WARD Christine E. Force–velocity relationships for isolated mouse soleus muscles during after-loaded and quick-released isotonic contractions. (PC) 161*P*

WATERLOW J.C. *see* JACKSON A.A., SLEVIN K. & WATERLOW J.C. A model of whole body protein turnover in man that includes a pool turning over by life-time kinetics. (C) 143*P*

WATSON M.S. *see* DEARNLEY D.P., GOMEZ S., HALL M., LUSH D., POPE R. & WATSON M.S. A computerized system for analysing lung function data in man. (D) 4*P*

WECKSTRÖM M. *see* LAUGHLIN S.B. & WECKSTRÖM M. The activation of a slow voltage-depedent potassium conductance is crucial for light adaptation in blowfly receptos. (PC) 200*P*

WEDLEY J.R. *see* GALLAGHER J., GARLICK P., HAMANN W., KEEVIL S., PETRICCIONE DI VADI P., PORTER D. & WEDLEY J.R. The use of magnetic resonance spectroscopy in assessing the efficacy of chemical lumbar sympathectomy in ischaemic vascular disease. (D) 3*P*

WELCH K.M.A. *see* BARKLEY G.L., GARDNER-MEDWIN A.R., MORAN J., NAGEL-LEIBY S., SIMKINS R.T., TEPLEY N. & WELCH K.M.A. Magnetic fields during spreading depression in the cortex of anaesthetized rabbits. (C) 101*P*

WELLER U. *see* DAUZENROTH M-E., DOLLY J.O., HABERMANN E., MOCHIDA S., POULAIN B., TAUC L., WADSWORTH J.D.F. & WELLER U. Light chain of tetanus toxin can be internalized into a cholinergic synapse of *Aplysia* by the heavy chain of botulinum neurotoxin and depresses the transmitter release. (C) 71*P*

WELSH Diana M. *see* NISTRI A., SIVILOTTI Lucia & WELSH Diana M. Effects of NMDA and its antagonists on excitatory synaptic transmission in the frog optic tectum *in vitro*. (PC) 151*P*

WERNER W. *see* BENNETT K.A., LEMON R.N. & WERNER W. Indirect excitation of corticospinal neurones by intracortical stimulation in the conscious monkey. (C) 103*P*

WEST J.M. *see* STEPHENSON D.G. & WEST J.M. Can the contractile apparatus in single muscle fibres from the extensor claw muscle of the yabby be fully activated by Sr^{2+} and Ca^{2+}? (PC) 162*P*

WEST N.H. *see* BACKHOUSE S.S., HARPER A.A., VAN VLIET B.N. & WEST N.H. Localization and conduction velocities of arterial baroreceptors in the anaesthetized snake. (C) 135*P*

WESTLING G. *see* BIGLAND-RITCHIE B., JOHANSSON R.S., THOMAS C.K. & WESTLING G. Twitch fusion in human motor units studied by motor axon stimulation. (C) 54*P*

WHITAKER Elaine M. *see* GUNASENA Karen T., MORRISON J.F.B., NIMMO A.J. & WHITAKER Elaine M. The effect of oestradiol and progesterone on neurokinin receptors in rat uterus. (C) 75*P*

WHITE D.C.S. *see* PECKHAM M., SPARROW J.C. & WHITE D.C.S. Mechanical responses of skinned flight muscle from mutants of *Drosophila melanogaster* with abnormal amounts of actin or myosin. (PC) 160*P*

WHITE J.M. *see* ALMERS W., IWATA A., SPRUCE A.E. & WHITE J.M. Patch clamp studies of fusion of human erythrocytes to murine fibroblasts mediated by influenza virus fusion protein. (C) 95*P*

WIDDAS W.F. *see* BAKER G.F. & WIDDAS W.F. Red cell volume compensation in isosmotic glucose. (C) 87*P*

WIER W.G. *see* BALKE C.W. & WIER W.G. Two types of calcium current in isolated guinea-pig ventricular myocytes: accurate separation by voltage-dependence of steady-state inactivation. (C) 31*P*

WIER W.G. *see* SIPIDO K.R. & WIER W.G. $[Ca^{2+}]_i$ transients in isolated guinea-pig ventricular myocytes in the absence of sodium–calcium (Na–Ca) exchange and sarcoplasmic reticulum function. (PC) 170*P*

WILLIAMS A.J. *see* SITSAPESAN R. & WILLIAMS A.J. Caffeine activation of single calcium-release channels from the sheep cardiac sarcoplasmic reticulum. (C) 26*P*

WILLIAMS A.J. *see* HOLMBERG S.R.M. & WILLIAMS A.J. Effects of doxorubicin and mitoxantrone on single calcium release channels from isolated sheep cardiac sarcoplasmic reticulum. (PC) 202*P*

WILLIAMS Pamela E. *see* GOLDSPINK Geoffrey & WILLIAMS Pamela E. The effect of immobilization on gene expression and post-natal determination of muscle fibre type in the mouse soleus muscle. (C) 52*P*

WILSON R.W. *see* TAYLOR E.W. & WILSON R.W. The cardiovascular and respiratory responses to intra-arterial injection of adrenaline in the dogfish *Scyliorhinus canicula* L. (C) 133*P*

WINLOW W. *see* SYED N.I. & WINLOW W. An identified, high-order neurone driving the whole animal withdrawal response in *Lymnaea*. (C) 72*P*

WINTERHAGER J.-M. *see* MACHEREY H.-J., PETERSEN K.-U., SPRAKTIES G. & WINTERHAGER J.-M. Effects of putative Cl channel blockers on electroneutral and electrogenic HCO_3 secretion by guinea-pig gallbladder epithelium *in vitro*. (C) 22*P*

WINTERTON S.J. *see* FLORES N.A., JUHASZ C., SHERIDAN D.J. & WINTERTON S.J. An IBM PC AT based system for the analysis of cardiac action potentials in real time. (D) 8*P*

WOAKES A.J. *see* ANDERSON L.S., BUTLER P.J., ROBERTS C., SMALE K., SNOW D.H. & WOAKES A.J. Partial pressures of arterial and end-tidal gases during graded exercise in thoroughbred race-horses. (C) 131*P*

WOOD J.N. *see* DOCHERTY R.J., FIEKERS J.F., WOOD J.N. & YEATS J.C. Histamine and prostaglandin D_2 modulate voltage-dependent calcium currents in adult rat sensory neurones in culture. (C) 36*P*

WOODWARD S.K.A. *see* DANTZIG J.A., GOLDMAN Y.E., LUTTMANN M.L., TRENTHAM D.R. & WOODWARD S.K.A. Binding of caged ATP diastereoisomers to rigor cross-bridges in glycerol-extracted fibres of rabbit psoas muscle. (C) 61*P*

WU Mei-Lin *see* VAUGHAN-JONES R.D. & WU Mei-Lin Intracellular intrinsic buffering power varies with pH_i in the isolated sheep cardiac Purkinje fibre. (PC) 165*P*

WYLLIE D.J.A. *see* CULL-CANDY S.G., MATHIE A., SYMONDS C.J. & WYLLIE D.J.A. Distribution of quisqualate and kainate receptors in rat type-2 astrocytes and their progenitor cells in culture. (PC) 195*P*

YANG J.F. *see* LLEWELLYN M., PROCHAZKA A. & YANG J.F. Human H-reflexes are reduced in difficult beam-walking compared to stance and normal walking. (C) 104*P*

YEATS J.C. *see* DOCHERTY R.J., FIEKERS J.F., WOOD J.N. & YEATS J.C. Histamine and prostaglandin D_2 modulate voltage-dependent calcium currents in adult rat sensory neurones in culture. (C) 36*P*

YEH Y. *see* BASKIN R.J., SHEN S. & YEH Y. Optical measurements on isolated single fibres from insect flight muscle. (C) 55*P*

YOSHINO M. *see* KAO C.Y., WAKUI M., WANG S.Y. & YOSHINO M. The outward current of the isolated rat myometrium. (C) 20*P*

YUDILEVICH D.L. *see* BARROS L.F., BUSTAMANTE J.C., JARVIS S.M. & YUDILEVICH D.L. Binding of nitrobenzylthioinosine, an inhibitor of facilitated-diffusion nucleoside transport, to brush-border and basal membrane vesicles isolated from the human placenta. (PC) 179*P*

YULE D.I. *see* DUNNE M.J., GALLACHER D.V., PETERSEN O.H. & YULE D.I. The electrogenic sodium–alanine co-transport system in RINm5F insulin-secreting cells: patch-clamp and single cell $[Ca^{2+}]_i$ measurements. (C) 96*P*

ZEIDEL M.L. *see* HARRIS H.W. Jr, JANOSHAZI A., KIKERI D., SOLOMON A.K. & ZEIDEL M.L. Proton permeability of toad bladder aggrephore membranes: role of the water channel in proton transport. (C) 145*P*

ZEITLIN I.J. *see* MOSHI M.J., PARRATT J.R., WAINWRIGHT C.L. & ZEITLIN I.J. Acid optimum kinin-forming enzymes in dog cardiovascular tissues. (PC) 177*P*

ZHOU Dan *see* ST JOHN Walter M. & ZHOU Dan Differing control of neural activities during various portions of expiration in the cat 189

ZORAN M.J. *see* HAYDON P.G., LUKOWIAK K., MAN-SON-HING H.J. & ZORAN M.J. Mechanisms of synaptic plasticity: modulation of secretory apparatus in cultured neuronal somata of *Helisoma trivolvis*. (C) 111*P*

ZORAN M.J. *see* HAYDON P.G., LUKOWIAK K., MAN-SON-HING H.J. & ZORAN M.J. Mechanisms of synaptic plasticity: Use of a giant soma–soma synapse in cultured neuronal somata of *Helisoma trivolvis*. (PC) 147*P*